

KEAP1 Protein, Human, Recombinant (His & GST & Avi)

General Information

Synonyms:	kelch like ECH associated protein 1; INRF2; KEAP-1; KLHL19
Protein Construction:	A DNA sequence encoding the human KEAP1 (Q14145) (Gln 2-Cys 624) was fused with the N-terminal polyhistidine-tagged GST tag at the N-terminus linked by the AVI tag. Predicted N terminal: Met
Species:	Human
Expression Host:	Baculovirus Insect Cells
Accession:	Q14145
Molecular Weight:	99.2 kDa (predicted); 85 kDa (reducing conditions)

QC Testing

Biological Activity:	Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 90 % as determined by SDS-PAGE
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing 20 mM Tris, 500 mM NaCl, 10% gly, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:
Reconstituted with sterile deionized water to 0.25 mg/mL. Reconstitution conditions may vary depending on the lot.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Kelch-like ECH-associated protein 1, also known as a cytosolic inhibitor of Nrf2, Kelch-like protein 19, KEAP1, and INRF2, is a cytoplasm and nucleus protein that contains one BACK (BTB/Kelch associated) domain, one BTB (POZ) domain, and six Kelch repeats. KEAP1 / INRF2 is broadly expressed, with the highest levels in skeletal muscle. KEAP1 / INRF2 is a key regulator of the NRF2 transcription factor, which transactivates the antioxidant response

element (ARE) and upregulates numerous proteins involved in antioxidant defense. Under basal conditions, KEAP1 / INRF2 targets NRF2 for ubiquitination and proteolytic degradation and as such is responsible for the rapid turnover of NRF2. KEAP1 / INRF2 retains NFE2L2 / NRF2 in the cytosol. KEAP1 / INRF2 functions as a substrate adapter protein for the E3 ubiquitin ligase complex formed by CUL3 and RBX1. It targets NFE2L2 / NRF2 for ubiquitination and degradation by the proteasome, thus resulting in the suppression of its transcriptional activity and the repression of antioxidant response element-mediated detoxifying enzyme gene expression. KEAP1 / INRF2 may also retain BPTF in the cytosol. It targets PGAM5 for ubiquitination and degradation by the proteasome.

Reference

- Strachan GD. et al., 2004, *Biochemistry*. 43: 12113-22.
Zhang DD. et al., 2004, *Mol Cell Biol*. 24: 10941-53.
Zhang DD. et al., 2003, *Mol Cell Biol*. 23: 8137-51.
Lo S-C. et al., 2006, *J Biol Chem*. 281: 37893-903.
Buckley BJ. et al., 2008, *Free Radic Biol Med*. 44 (4) :692-8.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481