

Kallikrein 3/KLK3 Protein, Human, Recombinant (His)

General Information

Synonyms:	APS;KLK2A1;PSA;kallikrein related peptidase 3;Hk3
Protein Construction:	Ala18-Pro261
Species:	Human
Expression Host:	HEK293 Cells
Accession:	P07288
Molecular Weight:	28.3 kDa (predicted); 32-38 kDa (reducing conditions)

QC Testing

Biological Activity:	Measured by its ability to cleave the colorimetric peptide substrate, Succinyl-Arg-Pro-Tyr-p-Nitroanilide(Suc-RPY-pNA). The specific activity is >100 pmoles/min/ μ g. (Activation description: The proenzyme needs to be activated by Thermolysin for an activated form)
Purity:	$\geq 95\%$ as determined by SDS-PAGE. $\geq 90\%$ as determined by SEC-HPLC.
Endotoxin:	< 1.0 EU/ μ g of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μ m filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μ g/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C . For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Kallikrein-related peptidase 3 (KLK3), also known as prostate-specific antigen (PSA), is the most useful biomarker for prostate cancer (PCa). KLK3 is suggested to play a role in regulating cancer growth through anti-angiogenic activity in vivo and in vitro.

Reference

Koistinen H, et al. (2008) Development of peptides specifically modulating the activity of KLK2 and KLK3. *Biol Chem.* 389(6): 633-42.

Yousef GM, et al. (2002) Kallikreins, steroid hormones and ovarian cancer: is there a link? *Minerva Endocrinol.* 27(3): 157-66.

Parikh H, et al. (2011) Fine mapping the KLK3 locus on chromosome 19q13.33 associated with prostate cancer susceptibility and PSA levels. *Hum Genet.* 129(6): 675-85.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481