

OX40/TNFRSF4 Protein, Mouse, Recombinant (hFc)

General Information

Synonyms:	TXGP1L;Txgp1;Ox40;CD134;ACT35;Ly-70;tumor necrosis factor receptor superfamily, member 4
Protein Construction:	A DNA sequence encoding the extracellular domain of mouse TNFRSF4 (NP_035789.1) (Met 1-Pro 211) was fused with the Fc region of human IgG1 at the C-terminus. Predicted N terminal: Val 20
Species:	Mouse
Expression Host:	HEK293 Cells
Accession:	B1ASL3
Molecular Weight:	48.3 kDa (predicted); 60 kDa (reducing condition, due to glycosylation)

QC Testing

Biological Activity:	Immobilized Recombinant Mouse OX40L/TNFSF4 Protein (His Tag) (Cat#TMPY-06018) at 0.5 µg/mL (100 µL/well) can bind Recombinant Mouse TNFRSF4/OX40/CD134 Protein (Fc Tag) (Cat#TMPY-02368), the EC50 is 0.8-2.5 ng/mL (QC tested).
Purity:	> 92 % as determined by SDS-PAGE
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:
A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

OX40 (CD134) and its binding partner, OX40L (CD252), are members of the tumor necrosis factor receptor/tumor necrosis factor superfamily, is known to break an existing state of tolerance in malignancies, leading to a

reactivation of antitumor immunity. The interaction between OX40 and OX40L plays an important role in antigen-specific T-cell expansion and survival. OX40 and OX40L also regulate cytokine production from T cells, antigen-presenting cells, natural killer cells, and natural killer T cells, and modulate cytokine receptor signaling. In line with these important modulatory functions, OX40-OX40L interactions have been found to play a central role in the development of multiple inflammatory and autoimmune diseases, making them attractive candidates for intervention in the clinic. Conversely, stimulating OX40 has shown it to be a candidate for therapeutic immunization strategies for cancer and infectious disease. Cancer Immunotherapy Co-stimulatory Immune Checkpoint Targets Immune Checkpoint Immune Checkpoint Detection: Antibodies Immune Checkpoint Detection: ELISA Antibodies Immune Checkpoint Detection: WB Antibodies Immune Checkpoint Proteins Immune Checkpoint Targets Immunotherapy Targeted Therapy

Reference

- Compaan D.M., et al. (2006) . The crystal structure of the costimulatory OX40-OX40L complex. Structure 14:1321-1330.
- Kawamata S., et al. (1998). Activation of OX40 signal transduction pathways leads to tumor necrosis factor receptor-associated factor (TRAF) 2- and TRAF5-mediated NF-kappaB activation. J. Biol. Chem. 273:5808-5814.
- Byun M., (2013) Inherited human OX40 deficiency underlying classic Kaposi sarcoma of childhood. J. Exp. Med. 210:1743-1759.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481