

## Nectin-2 Protein, Cynomolgus, Recombinant (His)

### General Information

Synonyms:	poliovirus receptor-related 2 (herpesvirus entry mediator B)
Protein Construction:	Gln32-Gly360
Species:	Cynomolgus
Expression Host:	HEK293 Cells
Accession:	A0A2K5U084
Molecular Weight:	36.53 kDa (predicted); 50-65 kDa (due to glycosylation)

### QC Testing

Biological Activity:	Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 95% as determined by Bis-Tris PAGE; > 95% as determined by HPLC
Endotoxin:	< 1.0 EU/ $\mu$ g of the protein as determined by the LAL method.
Formulation:	Lyophilized from 0.22 $\mu$ m filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

### Preparation and Storage

#### Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100  $\mu$ g/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

#### Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

#### Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

### Protein Background

Cluster of Differentiation 112 (CD112), also known as poliovirus receptor related protein 2 (PVRL2 or PRR2), is a single-pass type I transmembrane glycoprotein belonging to the Immunoglobulin superfamily. CD112 protein also serves as an entry for certain mutant strains of herpes simplex virus and pseudorabies virus, and thus is involved in cell to cell spreading of these viruses. CD112 protein has been identified as the ligand for DNAM-1 (CD226), and the interaction of CD226/CD112 protein can induce NK cell- and CD8+T cell-mediated cytotoxicity and cytokine

secretion. CD112 has been regarded as a critical component in allergic reactions, and accordingly may function as a novel target for anti-allergic therapy.

### Reference

Bachelet I, et al. (2006) Mast cell costimulation by CD226/CD112 (DNAM-1/Nectin-2): a novel interface in the allergic process. *J Biol Chem.* 281(37): 27190-6.

Wang L, et al. (2009) Molecular cloning, characterization and three-dimensional modeling of porcine nectin-2/CD112. *Vet Immunol Immunopathol.* 132(2-4): 257-63.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481