

Prostatic Acid Phosphatase Protein, Mouse, Recombinant (His)

General Information

Synonyms:	5'-NT;9.104100899-104272570.1;Ppal;A030005E02Rik;PAP;Lap;acid phosphatase, prostate; FRAP
Protein Construction:	A DNA sequence encoding the mouse ACPD isoform 1 (Q8CE08-1) (Met1-Arg 381) was expressed, with a C-terminal polyhistidine tag. Predicted N terminal: Lys 32
Species:	Mouse
Expression Host:	HEK293 Cells
Accession:	Q8CE08-1
Molecular Weight:	42 kDa (predicted); 47 kDa (reducing condition, due to glycosylation)

QC Testing

Biological Activity:	Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 98 % as determined by SDS-PAGE
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Prostatic acid phosphatase (PAP, or ACPD), also known as prostatic specific acid phosphatase (PSAP), is an enzyme produced by the prostate. As a non-specific phosphomonoesterase, Prostatic acid phosphatase is synthesized and secreted into seminal plasma under androgenic control. The enzyme is a dimer of molecular weight around 100 kDa. Prostatic acid phosphatase is a clinically important protein for its relevance as a biomarker of prostate

carcinoma. Furthermore, it has a potential role in fertilization. The major action of PAP is to dephosphorylate macromolecules with the help of catalytic residues (His(12) and Asp(258)) that are located in the cleft between two domains. Cellular prostatic acid phosphatase (cPAP), an authentic tyrosine phosphatase, is proposed to function as a negative growth regulator of prostate cancer (PCa) cells in part through its dephosphorylation of ErbB-2. cPAP functions as a neutral protein tyrosine phosphatase (PTP) in prostate cancer cells and dephosphorylates HER-2/ErbB-2/Neu (HER-2: human epidermal growth factor receptor-2) at the phosphotyrosine (p-Tyr) residues. Injection of the secretory isoform of PAP has potent antinociceptive effects in mouse models of chronic pain. This enzyme exhibits ecto-5'-nucleotidase activity, is widely distributed, and implicated in the formation of chronic pain. Additionally, PAP could be a target molecule in specific immunotherapy for patients with nonprostate adenocarcinomas including colon and gastric cancers.

Reference

- Hassan MI, et al. (2010) Structural and functional analysis of human prostatic acid phosphatase. *Expert Rev Anticancer Ther.* 10(7): 1055-68.
- Chuang TD, et al. (2010) Human prostatic acid phosphatase, an authentic tyrosine phosphatase, dephosphorylates ErbB-2 and regulates prostate cancer cell growth. *J Biol Chem.* 285(31): 23598-606.
- Larsen RS, et al. (2009) A high throughput assay to identify small molecule modulators of prostatic acid phosphatase. *Curr Chem Genomics.* 3: 42-9.
- Zimmermann H. (2009) Prostatic acid phosphatase, a neglected ectonucleotidase. *Purinergic Signal.* 5(3): 273-5.
- Wang Y, et al. (2005) Prostatic acid phosphatase as a target molecule in specific immunotherapy for patients with nonprostate adenocarcinoma. *J Immunother.* 28(6): 535-41.
- Veeramani S, et al. (2005) Cellular prostatic acid phosphatase: a protein tyrosine phosphatase involved in androgen-independent proliferation of prostate cancer. *Endocr Relat Cancer.* 12(4): 805-22.
- Ostrowski WS, et al. (1994) Human prostatic acid phosphatase: selected properties and practical applications. *Clin Chim Acta.* 226(2): 121-9.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481