

ANGPT4/Angiopoietin-4 Protein, Human, Recombinant (hFc)

General Information

Synonyms:	angiopoietin 4;ANG4;ANG-3;AGP4
Protein Construction:	A DNA sequence encoding the human ANGPT4 (Q9Y264) C-terminal fragment (Met 282-Ile 503) was fused with the Fc region of human IgG1 at the N-terminus. Predicted N terminal: Glu
Species:	Human
Expression Host:	HEK293 Cells
Accession:	Q9Y264
Molecular Weight:	53.7 kDa (predicted); 66 kDa (reducing condition, due to glycosylation)

QC Testing

Biological Activity:	Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 90 % as determined by SDS-PAGE
Endotoxin:	< 1.0 EU/ μ g of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μ m filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:
A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:
It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:
In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Reference

Grosios K, et al. (1999) Assignment of ANGPT4, ANGPT1, and ANGPT2 encoding angiopoietins 4, 1 and 2 to human chromosome bands 20p13, 8q22.3?q23 and 8p23.1, respectively, by in situ hybridization and radiation hybrid mapping. *Cytogenet. Cell Genet.* 84 (1-2): 118-20.

Yamakawa M, et al. (2004) Expression of angiopoietins in renal epithelial and clear cell carcinoma cells: regulation by hypoxia and participation in angiogenesis. *Am J Physiol Renal Physiol.* 287 (4): 649-57.

Oliner J, et al. (2004) Suppression of angiogenesis and tumor growth by selective inhibition of angiopoietin-2. *Cancer Cell.* 6 (5): 507-16.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481