

EPCR Protein, Human, Recombinant (His)

General Information

| | |
|-----------------------|--|
| Synonyms: | EPCR;CCD41;CCCA;protein C receptor, endothelial |
| Protein Construction: | A DNA sequence encoding the extracellular domain of human PROCR (Q9UNN8) (Met 1-Thr 209), was fused with a polyhistidine tag at the C-terminus. Predicted N terminal: Ser 18 |
| Species: | Human |
| Expression Host: | HEK293 Cells |
| Accession: | Q9UNN8 |
| Molecular Weight: | 23.4 kDa (predicted); 37 kDa (reducing condition, due to glycosylation) |

QC Testing

| | |
|----------------------|--|
| Biological Activity: | Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first. |
| Purity: | ≥ 97 % as determined by SDS-PAGE. ≥ 90 % as determined by SEC-HPLC. |
| Endotoxin: | < 1.0 EU/μg of the protein as determined by the LAL method. |
| Formulation: | Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization. |

Preparation and Storage

Reconstitution:
A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Endothelial protein C receptor (EPCR), also known as activated protein C receptor (APC receptor) or PROCR, is a receptor for Protein C. Protein C plays an important role in many metabolism processes in humans and other animals after activated by binding to Endothelial protein C receptor (EPCR). Because of the EPCR is found primarily on endothelial cells (cells on the inside of blood vessels), activated protein C is found mainly near endothelial cells. Protein C is pleiotropic, with two main functions: anticoagulation and cytoprotection. Which function will be

performed depending on whether or not protein C remains bind to EPCR after activated. The anticoagulation occurs when it does not. In this case, protein C functions as an anticoagulant by irreversibly proteolytically inactivating Factor Va and Factor VIIIa, turning them into Factor Vi and Factor VIIIi respectively. When still bound to EPCR, activated protein C performs its cytoprotective effects, acting on the effector substrate PAR-1, protease-activated receptor-1. To a degree, APC's anticoagulant properties are independent of its cytoprotective ones, in that expression of one pathway is not affected by the existence of the other.

Reference

- Nicolaes GA,et al. (2003). Congenital and acquired activated protein C resistance. *Semin Vasc Med.* 3 (1): 33-46.
Esmon CT. (2003). The protein C pathway. *Chest* 124 (3): 26-32.
Mosnier LO,et al. (2007)The cytoprotective protein C pathway. *Blood.* 109: 3161-72.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481