

## IgG3 Protein, Human, Recombinant

### General Information

Synonyms:	IGHG3
Protein Construction:	A DNA sequence encoding the human IgG3 Fc region (P01860) (Glu 99-Lys 377) was expressed and purified. Predicted N terminal: Glu 99
Species:	Human
Expression Host:	HEK293 Cells
Accession:	P01860
Molecular Weight:	31.2 kDa (predicted); 38 kDa (reducing condition, due to glycosylation)

### QC Testing

Biological Activity:	Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	≥ 96 % as determined by SDS-PAGE. ≥ 95 % as determined by SEC-HPLC.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

### Preparation and Storage

**Reconstitution:**  
A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

**Stability & Storage:**

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

**Shipping:**

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

### Protein Background

IGHG3 (Immunoglobulin Heavy Constant Gamma 3 (G3m Marker), also known as IgG3) is a Protein Coding gene. Ig gamma-3 chain C region is a protein that in humans is encoded by the IGHG3 gene. Immunoglobulins, also known as antibodies, are membrane-bound or secreted glycoproteins produced by B lymphocytes. In the recognition phase of humoral immunity, the membrane-bound immunoglobulins serve as receptors which, upon binding of a specific antigen, trigger the clonal expansion and differentiation of B lymphocytes into immunoglobulins-

secreting plasma cells. Murine immunoglobulin G (IgG) plays an important role in mediating protective immune responses to malaria. Diseases associated with IGHG3 include Heavy Chain Disease and Gamma Heavy Chain Disease. Among its related pathways are IL4-mediated signaling events and the Creation of C4 and C2 activators.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481