

Nicastrin Protein, Mouse, Recombinant (hFc)

General Information

Synonyms:	Aph2;AA727311;Nct;NCSTN;nicastrin;mKIAA0253;D1Dau13e;9430068N19Rik;Kiaa0253
Protein Construction:	A DNA sequence encoding the mouse NCSTN (P57716) (Met1-Glu668) was expressed, fused with the Fc region of human IgG1 at the C-terminus. Predicted N terminal: Gly 28
Species:	Mouse
Expression Host:	HEK293 Cells
Accession:	P57716
Molecular Weight:	98.5 kDa (predicted); 117 kDa (reducing condition, due to glycosylation)

QC Testing

Biological Activity:	Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 95 % as determined by SDS-PAGE
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Nicastrin (NCST, or NCT), a single-pass membrane glycoprotein that harbors a large extracellular domain, is an essential component of the gamma-secretase complex. Several lines of evidence indicate that the members of these complexes could also contribute to the control of cell death. NCT controls cell death via phosphoinositide 3-kinase/Akt and p53-dependent pathways and that this function remains independent of the activity and molecular integrity of the gamma-secretase complexes. Increasing pieces of evidence have shown that Nicastrin/NCSTN

plays a crucial role in gamma-cleavage of the amyloid precursor protein (APP). The glycoprotein Nicastrin is an essential component of the gamma-secretase complex, a high molecular weight complex that also contains the presenilin proteins, Aph-1 and Pen-2. The gamma-secretase complex is not only involved in APP processing but also in the processing of an increasing number of another type I integral membrane proteins. As the largest subunit of the gamma-secretase complex, Nicastrin plays a crucial role in its activation. Inhibition of NCSTN demonstrated an altered gamma-cleavage activity, suggesting its potential implication in developing Alzheimer's disease (AD). Besides, Nicastrin can function to maintain epithelial to mesenchymal transition during breast cancer progression. Anti-nicastrin polyclonal and monoclonal antibodies were able to decrease notch1 and vimentin expression and reduced the invasive capacity of breast cancer cells in vitro.

Reference

- He G, et al. (2007) Degradation of nicastrin involves both proteasome and lysosome. *J Neurochem.* 101(4): 982-92.
- Ma Z, et al. (2009) Association between promoter polymorphisms of the nicastrin gene and sporadic Alzheimer's disease in North Chinese Han population. *Neurosci Lett.* 458(3): 136-9.
- Hayashi I, et al. (2009) Single chain variable fragment against nicastrin inhibits the gamma-secretase activity. *J Biol Chem.* 284(41): 27838-47.
- Pardossi-Piquard R, et al. (2009) p53-dependent control of cell death by nicastrin: lack of requirement for presenilin-dependent gamma-secretase complex. *J Neurochem.* 109(1): 225-37.
- Filipovi? A, et al. (2011) Biological and clinical implications of nicastrin expression in invasive breast cancer. *Breast Cancer Res Treat.* 125(1): 43-53.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481