

TGFBR2 Protein, Rhesus, Recombinant (hFc)

General Information

| | |
|-----------------------|---|
| Synonyms: | transforming growth factor, β receptor II; transforming growth factor, beta receptor II (70/80kDa) |
| Protein Construction: | A DNA sequence encoding the rhesus TGFBR2 (NP_001248080.1) (Met1-Asp159) was expressed with the Fc region of human IgG1 at the C-terminus. Predicted N terminal: Ile 24 |
| Species: | Rhesus |
| Expression Host: | HEK293 Cells |
| Accession: | H9YUL0 |
| Molecular Weight: | 42.4 kDa (predicted) |

QC Testing

| | |
|----------------------|---|
| Biological Activity: | 1. Measured by its binding ability in a functional ELISA. 2. Immobilized Rhesus TGFBR2-His at 10 $\mu\text{g}/\text{mL}$ (100 $\mu\text{L}/\text{well}$) can bind Rhesus TGFBR2-Fc. The EC50 of Rhesus TGFBR2-Fc is 28-56 ng/mL. |
| Purity: | > 95 % as determined by SDS-PAGE |
| Endotoxin: | < 1.0 EU/ μg of the protein as determined by the LAL method. |
| Formulation: | Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization. |

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C . For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

TGFBR2 is a member of the Ser/Thr protein kinase family and the TGFBR receptor subfamily. It is a transmembrane protein. TGFBR2 is comprised of a C-terminal protein kinase domain and an N-terminal ectodomain. The ectodomain consists of a compact fold containing nine beta-strands and a single helix stabilized by a network of

six intra strand disulfide bonds. The folding topology includes a central five-stranded antiparallel beta-sheet, eight-residues long at its centre, covered by a second layer consisting of two segments of two-stranded antiparallel beta-sheets. TGFBR2 has a protein kinase domain, forms a heterodimeric complex with another receptor protein, and binds TGF-beta. This receptor/ligand complex phosphorylates proteins, which then enter the nucleus and regulate the transcription of a subset of genes related to cell proliferation. Mutations in TGFBR2 gene have been associated with Marfan syndrome, Loeys-Deitz Aortic Aneurysm Syndrome, and the development of various types of tumors. TGFBR2 attenuates the biological activities of TGF-beta in colorectal cancer. TGFBR2 expression is increased in oral squamous cell carcinoma cells. Its expression is decreased by IL-1beta while inducing Sp3 via NFkappaB. TGFBR2 and TGFBR1 are involved in the antiestrogenic activity.

Reference

- Yu Y, et al. (2012) MicroRNA-21 induces stemness by downregulating transforming growth factor beta receptor 2 (TGFBR2) in colon cancer cells. *Carcinogenesis*. 33(1):68-76.
- Shima K, et al. (2011) TGFBR2 and BAX mononucleotide tract mutations, microsatellite instability, and prognosis in 1072 colorectal cancers. *PLoS One*. 6(9):e25062.
- Biros E, et al. (2011) Meta-analysis of the association between single nucleotide polymorphisms in TGF-beta receptor genes and abdominal aortic aneurysm. *Atherosclerosis*. 219(1):218-23.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481