

## IL-6 Protein, Rhesus, Recombinant

### General Information

Synonyms:	IL6;interleukin 6;IL-6
Protein Construction:	A DNA sequence encoding the the rhesus IL6 (NP_001036198.2 ) (Pro29-Met212) was expressed and purified. Predicted N terminal: Met
Species:	Rhesus
Expression Host:	E. coli
Accession:	A0A1D5QRZ4
Molecular Weight:	21 kDa (predicted)

### QC Testing

Biological Activity:	Measured in a cell proliferation assay using TF-1 human erythroleukemic cells. The ED50 for this effect is typically 5-30 ng/ml.
Purity:	> 85 % as determined by SDS-PAGE
Endotoxin:	Please contact us for more information.
Formulation:	Lyophilized from a solution filtered through a 0.22 $\mu$ m filter, containing 50% acetonitrile, 0.1% TFA. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

### Preparation and Storage

**Reconstitution:**  
Reconstituted with sterile deionized water to 0.25 mg/mL. Reconstitution conditions may vary depending on the lot.

**Stability & Storage:**

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

**Shipping:**

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

### Protein Background

Interleukin-6 (IL-6) is a multifunctional  $\alpha$ -helical cytokine that regulates cell growth and differentiation of various tissues, which is known particularly for its role in the immune response and acute phase reactions. IL-6 protein is secreted by a variety of cell types including T cells and macrophages as a phosphorylated and variably glycosylated molecule. It exerts actions through its heterodimeric receptor composed of IL-6R that lacks the tyrosine/kinase domain and binds IL-6 with low affinity, and ubiquitously expressed glycoprotein 130 (gp130) that

binds the IL-6. IL-6R complex with high affinity and thus transduces signals. IL-6 is also involved in hematopoiesis, bone metabolism, and cancer progression, and has been defined as an essential role in directing the transition from innate to acquired immunity. Cancer Immunotherapy/Immune Checkpoint Immunotherapy/Targeted Therapy

### Reference

Heinrich PC, et al. (2003). Principles of interleukin-6-type cytokine signalling and its regulation. *Biochem J.* 374: 1-20.

Rose-John S, et al. (2007) The IL-6/sIL-6R complex as a novel target for therapeutic approaches. *Expert Opin Ther Targets.* 11(5): 613-24.

Dinh W, et al. (2009) Elevated plasma levels of TNF-alpha and interleukin-6 in patients with diastolic dysfunction and glucose metabolism disorders. *Cardiovasc Diabetol.* 8:58.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E\_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481