

## EBAG9 Protein, Human, Recombinant (His)

### General Information

Synonyms:	EB9;PDAF;estrogen receptor binding site associated, antigen, 9
Protein Construction:	A DNA sequence encoding the mature form of human EBAG9 (O00559) (Arg28-Ser213) was expressed with a polyhistidine tag at the N-terminus. Predicted N terminal: His
Species:	Human
Expression Host:	E. coli
Accession:	O00559
Molecular Weight:	23 kDa (predicted); 33 kDa (reducing conditions)

### QC Testing

Biological Activity:	Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 90 % as determined by SDS-PAGE
Endotoxin:	Please contact us for more information.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

### Preparation and Storage

**Reconstitution:**  
A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

**Stability & Storage:**  
It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

**Shipping:**  
In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

### Protein Background

RCAS1, also known as EBAG9, is a tumor-associated antigen that is expressed at high frequency in a variety of cancers. RCAS1 gene was identified as an estrogen-responsive gene. Regulation of transcription by estrogen is mediated by estrogen receptor which binds to the estrogen-responsive element (ERE) found in the 5'-flanking region of RCAS1 gene. Two transcript variants differing in the 5' UTR, but encoding the same protein, have been identified for RCAS1 gene. EBAG9 may participate in suppression of cell proliferation and induces apoptotic cell

death through activation of interleukin-1-beta converting enzyme (ICE)-like proteases.

Reference

Ohshima K. et al., 2001, Clin Exp Immunol. 123 (3): 481-6.

Ikeda K. et al., 2000, Biochem Biophys Res Commun. 273 (2): 654-60.

Nakashima M. et al., 1999, Nat Med. 5 (8): 938-42.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481