

MERS-CoV Spike Protein (S1+S2 ECD, aa 1-1297, His)

General Information

Synonyms:	coronavirus spike Protein, MERS-CoV;ncov spike Protein, MERS-CoV;coronavirus s1 Protein, MERS-CoV;ncov RBD Protein, MERS-CoV;cov spike Protein, MERS-CoV;ncov s2 Protein, MERS-CoV;coronavirus s2 Protein, MERS-CoV;RBD Prote;ncov s1 Protein, MERS-CoV
Protein Construction:	A DNA sequence encoding the MERS-CoV extracellular domain of spike protein (AFS88936.1) (Met1-Trp1297) was fused with a polyhistidine tag at the C-terminus. Predicted N terminal: Tyr 18
Species:	MERS-CoV
Expression Host:	Baculovirus Insect Cells
Accession:	K0BRG7
Molecular Weight:	44.10 kDa (predicted)

QC Testing

Biological Activity:	Measured by its binding ability in a functional ELISA.1. Immobilized Sh(2C-EMC)(1-1297) at 10 µg/mL (100 µl/well) can bind biotinylated DPP4 , the EC50 of can biotinylated DPP4 is 0.02-0.04 ug/mL.2. Immobilized Sh(2C-EMC)(1-1297) at 10 µg/mL (100 µl/well) can bind biotinylated Fc-DPP4 , the EC50 of can biotinylated Fc-DPP4 is 0.01-0.02 ug/mL.
Purity:	> 85 % as determined by SDS-PAGE
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing 20 mM Tris, 300 mM NaCl, 10% glycerol, pH7.5. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:	Reconstituted with sterile deionized water to 0.25 mg/mL. Reconstitution conditions may vary depending on the lot.
Stability & Storage:	It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots. <small>Actual storage temperature shall be subject to the COA.</small>
Shipping:	In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

The spike (S) glycoprotein of coronaviruses contains protrusions that will only bind to certain receptors on the host cell. Known receptors bind S1 are ACE2, angiotensin-converting enzyme 2; DPP4, dipeptidyl peptidase-4; APN, aminopeptidase N; CEACAM, carcinoembryonic antigen-related cell adhesion molecule 1; Sia, sialic acid; O-ac Sia, O-acetylated sialic acid. The spike is essential for both host specificity and viral infectivity. The term 'peplomer' is typically used to refer to a grouping of heterologous proteins on the virus surface that function together. The spike (S) glycoprotein of coronaviruses is known to be essential in the binding of the virus to the host cell at the advent of the infection process. It's been reported that SARS-CoV-2 (COVID-19 coronavirus, 2019-nCoV) can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity. The main functions for the Spike protein are summarized as: Mediate receptor binding and membrane fusion; Defines the range of the hosts and specificity of the virus; Main component to bind with the neutralizing antibody; Key target for vaccine design; Can be transmitted between different hosts through gene recombination or mutation of the receptor binding domain (RBD), leading to a higher mortality rate.

Reference

Xiao X, et al. (2004) The SARS-CoV S glycoprotein. *Cell Mol Life Sci.* 61 (19-20): 2428-30.

Shen S, et al. (2007) Expression, glycosylation, and modification of the spike (S) glycoprotein of SARS CoV. *Methods Mol Biol.* 379: 127-35.

Du L, et al. (2009) The spike protein of SARS-CoV--a target for vaccine and therapeutic development. *Nat Rev Microbiol.* 7 (3): 226-36.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481