

SLAM/CD150 Protein, Rat, Recombinant (His)

General Information

Synonyms:	signaling lymphocytic activation molecule family member 1
Protein Construction:	A DNA sequence encoding the rat SLAMF1 (D3ZAD7) (Met1-Leu242) was expressed with a polyhistidine tag at the C-terminus. Predicted N terminal: Thr 25
Species:	Rat
Expression Host:	HEK293 Cells
Accession:	D3ZAD7
Molecular Weight:	17.4 kDa (predicted); 44-47 kDa (reducing condition, due to glycosylation)

QC Testing

Biological Activity:	Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 95 % as determined by SDS-PAGE
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:
A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

CD150/signaling lymphocytic activation molecule (SLAM) is a cell surface sialylated phosphoglycoprotein and belongs to the CD2 subset of the Ig superfamily of type I transmembrane glycoproteins. The CD150 receptor is expressed on thymocytes, activated and memory T cells, B cells, platelets, natural killer T cells, and mature dendritic cells, and is also detected on tumor cells of Hodgkin's lymphoma (HL) and diffuse large B-cell lymphoma with an activated B cell phenotype. Additionally, it is the immune cell receptor for measles virus (MV). As a self-

ligand, CD150 performs diverse immunologic functions including T/B-cell costimulation, induction of interferon γ (IFN- γ) in Th1 T-cell clones, redirection of Th2 clones to a Th1 or Th0 phenotype, and inhibition of apoptosis in B cells. Furthermore, CD150 was shown to be the second receptor for measles virus in addition to CD46, and the distribution of SLAM on various cell lines is consistent with their susceptibility to clinical isolates of measles virus.

Reference

Tatsuo H, et al. (2002) The morbillivirus receptor SLAM (CD150). *Microbiol Immunol.* 46(3): 135-42.

Sidorenko SP, et al. (2003) The dual-function CD150 receptor subfamily: the viral attraction. *Nat Immunol.* 4(1): 19-24.

Yurchenko MY, et al. (2010) CD150 regulates JNK1/2 activation in normal and Hodgkin's lymphoma B cells. *Immunol Cell Biol.* 88(5): 565-74.

Leonard VH, et al. (2010) Measles virus selectively blind to signaling lymphocytic activation molecule (SLAM ; CD150) is attenuated and induces strong adaptive immune responses in rhesus monkeys. *J Virol.* 84(7): 3413-20.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481