

TSPAN8 Protein, Human, Recombinant (mFc)

General Information

Synonyms:	TM4SF3;tetraspanin 8;CO-029
Protein Construction:	A DNA sequence encoding the human TSPAN8 (P19075) (Lys110-Asn205) was expressed with the Fc region of mouse IgG1 at the N-terminus. Predicted N terminal: Asp
Species:	Human
Expression Host:	HEK293 Cells
Accession:	P19075
Molecular Weight:	58 kDa (predicted); 46 kDa (reducing condition, due to glycosylation)

QC Testing

Biological Activity:	Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 95 % as determined by SDS-PAGE
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:
A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:
It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:
In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Tetraspanin 8 (TSPAN8) as an important modulator of melanoma invasiveness, and several of its transcriptional regulators, which affect TSPAN8 expression during melanoma progression toward an invasive stage. p53 as a negative regulator of Tspan8 expression. p53 as a regulator of melanoma invasion and the concept that reactivating p53 could provide a strategy for modulating not only proliferative but also invasive capacity in melanoma treatment. Tetraspanin 8 (TSPAN8) is a tumor-associated antigen implicated in tumor progression and

metastasis. TSPAN8 may play an important role in mCRC cell invasion. TSPAN8 was overexpressed in human gastric cancer tissues and gastric cancer cell lines compared with the normal. TSPAN8 overexpression promoted cell proliferation and invasion, while TSPAN8 suppression inhibited cell proliferation and invasion. TSPAN8 could activate the ERK MAPK pathway in gastric cancer cells, and MEK-ERK inhibition reversed the effects of TSPAN8 overexpression on cell proliferation and invasion.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481