

EDA2R Protein, Cynomolgus, Recombinant (hFc)

General Information

Synonyms:	ectodysplasin A2 receptor
Protein Construction:	A DNA sequence encoding the cynomolgus EDA2R (NP_001180396.1) (Met1-Glu136) was expressed with the Fc region of human IgG1 at the C-terminus. Predicted N terminal: Met
Species:	Cynomolgus
Expression Host:	HEK293 Cells
Accession:	A0A5F8A4K0
Molecular Weight:	38.4 kDa (predicted); 116, 58 and 53 kDa (reducing conditions)

QC Testing

Biological Activity:	Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 90 % as determined by SDS-PAGE
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:
A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:
It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:
In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Tumor necrosis factor receptor superfamily member 27, also known as X-linked ectodysplasin-A2 receptor, EDA-A2 receptor, EDA2R, XEDAR and TNFRSF27, is a single-pass type III membrane protein. TNFRSF27 / EDA2R contains three TNFR-Cys repeats. It is a new member of the tumor necrosis factor receptor family that has been shown to be highly expressed in ectodermal derivatives during embryonic development and binds to ectodysplasin-A2 (EDA-A2). TNFRSF27 / EDA2R is a receptor for EDA isoform A2, but not for EDA isoform A1.

A DRUG SCREENING EXPERT

TNFRSF27 / EDA2R mediates the activation of the NF-kappa-B and JNK pathways. The activation seems to be mediated by binding to TRAF3 and TRAF6. Ectodysplasin, a member of the tumor necrosis factor family, is encoded by the anhidrotic ectodermal dysplasia EDA gene. Mutations in EDA give rise to a clinical syndrome characterized by loss of hair, sweat glands, and teeth. EDA-A1 and EDA-A2 are two isoforms of ectodysplasin that differ only by an insertion of two amino acids. This insertion functions to determine receptor binding specificity, such that EDA-A1 binds only the receptor EDAR, whereas EDA-A2 binds only the related, but distinct, X-linked ectodysplasin-A2 receptor (XEDAR).

Reference

Yan M., et al., 2000, Science. 290: 523-527.

Sinha S.K., et al., 2002, J. Biol. Chem. 277: 44953-44961.

Prodi, D.A. et al., 2008, J Invest Dermatol 128 (9):2268-70.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481