

## OX40L/TNFSF4 Protein, Cynomolgus, Recombinant (mFc)

### General Information

Synonyms:	tumor necrosis factor (ligand) superfamily, member 4
Protein Construction:	A DNA sequence encoding the cynomolgus TNFSF4 (F7FL80) (Gln51-Leu183) was expressed with the Fc region of mouse IgG1 at the N-terminus. Predicted N terminal: Asp
Species:	Cynomolgus
Expression Host:	HEK293 Cells
Accession:	F7FL80
Molecular Weight:	42 kDa (predicted); 48 kDa (reducing conditions)

### QC Testing

Biological Activity:	Immobilized Cynomolgus mFc-TNFSF4 at 10 µg/ml (100 µl/well) can bind human TNFRSF4-Fch, The EC50 of human TNFRSF4-Fch is 0.23-0.55 µg/ml.
Purity:	> 85 % as determined by SDS-PAGE
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

### Preparation and Storage

#### Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

#### Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

#### Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

### Protein Background

OX-40L, also known as TNFSF4 and CD252, is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. OX-40L is an important costimulatory molecule that plays a crucial role in the regulation of T-cell-mediated immunity. The interaction of TNFSF4-TNFSF4 is involved in the pathogenesis of multiple autoimmune and inflammatory diseases such as systemic lupus erythematosus (SLE), carotid artery disease and cancer. OX-40L is a ligand for receptor TNFRSF4/OX40. It is found to play a role in T cell antigen-presenting cell (APC) interactions. In

surface Ig- and CD4-stimulated B cells, this cytokine along with CD7 has been shown to provide CD28-independent costimulatory signals to T cells. This protein and its receptor are reported to directly mediate adhesion of activated T cells to vascular endothelial cells. Cancer Immunotherapy Co-stimulatory Immune Checkpoint Targets Immune Checkpoint Immune Checkpoint Targets Immunotherapy Targeted Therapy

### Reference

Lei W. et al., 2012, Ann Acad Med Singapore. 41 (5): 200-4.

Lee YH. et al., 2012, Hum Immunol. 73 (10): 1050-4.

Weiguang Y. et al., 2012, PLoS One. 7 (8): e41277.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E\_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481