

MCP-3/CCL7 Protein, Mouse, Recombinant (His)

General Information

Synonyms:	chemokine (C-C motif) ligand 7;mcp3;marc;fic;Scya7;MCP-3
Protein Construction:	A DNA sequence encoding the mouse Ccl7 (NP_038682.1) (Gln24-Pro97) was expressed with a polyhistidine tag at the N-terminus. Predicted N terminal: Met
Species:	Mouse
Expression Host:	E. coli
Accession:	A9Z1Z1
Molecular Weight:	10.7 kDa (predicted)

QC Testing

Biological Activity:	Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 90 % as determined by SDS-PAGE.
Endotoxin:	Please contact us for more information.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS, 10% glycerol. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:	A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.
Stability & Storage:	It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots. <small>Actual storage temperature shall be subject to the COA.</small>
Shipping:	In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Chemokines are a family of small chemotactic cytokines, or proteins secreted by cells. Chemokines share the same structure similarities such as small size, and the presence of four cysteine residues in conserved locations in order to form their 3-dimensional shape. Some of the chemokines are considered pro-inflammatory which can be induced to recruit cells of the immune system to a site of infection during an immune response, while others are considered homeostatic and are implied in controlling the migration of cells during normal processes of tissue

maintenance and development. There are four members of the chemokine family: C-C chemokines, C chemokines, CXC chemokines and CX3C chemokines. The C-C chemokines have two cysteines nearby the amino terminus. There have been at least 27 distinct members of this subgroup reported for mammals, called C-C chemokine ligands-1 to 28. Chemokine ligand 7(CCL7), also known as MCP-3, is a member of the C-C chemokine subfamily of the chemokine family which is produced by certain tumor cells and by macrophages. It also owns two adjacent N-terminal cysteine residues. Chemokine ligand 7(CCL7) specifically attracts monocytes, and regulates macrophage function.

Reference

- Laing KJ, et al. (2004) Chemokines. *Developmental and comparative immunology*. 28 (5): 443-60.
- Cocchi F, et al. (1995) Identification of RANTES, MIP-1a, and MIP-1b as the major HIV-suppressive factor produced by CD8+T cells. *Science*. 270 (5243): 1811-5.
- Hori T, et al. (2008) CCL8 is a potential molecular candidate for the diagnosis of graft-versus-host disease. *Blood*. 111 (8): 4403-12.
- Biber K, et al. (2003) Expression of L-CCR in HEK 293 cells reveals functional responses to CCL2, CCL5, CCL7, and CCL8. *Journal of Leukocyte Biology*. 74 (2): 243-51.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481