

IGF1R/CD221 Protein, Human, Recombinant (His & GST)

General Information

Synonyms:	JTK13;insulin-like growth factor 1 receptor;IGF1 Receptor;CD221;IGFR;IGFIR;IGF-I R
Protein Construction:	A DNA sequence encoding the human IGF1R (NP_000866.1) cytoplasmic domain (Met 954-Cys 1367) was fused with the N-terminal polyhistidine-tagged GST tag at the N-terminus. Predicted N terminal: Met
Species:	Human
Expression Host:	Baculovirus Insect Cells
Accession:	P08069
Molecular Weight:	74.6 kDa (predicted)

QC Testing

Biological Activity:	The specific activity was determined to be >100 nmol/min/mg using Poly(Glu,Tyr) 4:1 as substrate.
Purity:	> 85 % as determined by SDS-PAGE
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Supplied as sterile 20 mM Tris, 500 mM NaCl, 20% glycerol, pH 7.4.

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store the product under sterile conditions at -20°C to -80°C. Samples are stable for up to 12 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

Proteins are shipped with blue ice.

Protein Background

The insulin-like growth factor-1 receptor (IGF1R) is a transmembrane tyrosine kinase involved in several biological processes including cell proliferation, differentiation, DNA repair, and cell survival. This a disulfide-linked heterotetrameric transmembrane protein consisting of two α and two β subunits, and among which, the α subunit is extracellular while the β subunit has an extracellular domain, a transmembrane domain, and a cytoplasmic tyrosine kinase domain. The IGF1R signaling pathway is activated in the mammalian nervous system from the early developmental stages. Its major effect on developing neural cells is to promote their growth and survival. This pathway can integrate its action with signaling pathways of growth and morphogenetic factors that induce

cell fate specification and selective expansion of specified neural cell subsets. Modulation of cell migration is another possible role that IGF1R activation may play in neurogenesis. In the mature brain, IGF-I binding sites have been found in different regions of the brain, and multiple reports confirmed a strong neuroprotective action of the IGF-IR against different pro-apoptotic insults. IGF1R is an important signaling molecule in cancer cells and plays an essential role in the establishment and maintenance of the transformed phenotype. Inhibition of IGF1R signaling thus appears to be a promising strategy to interfere with the growth and survival of cancer cells. IGF1R is frequently overexpressed by tumors and mediates proliferation and apoptosis protection. IGF signaling also influences hypoxia signaling, protease secretion, tumor cell motility, and adhesion, and thus can affect the propensity for invasion and metastasis. Therefore, IGF1R is now an attractive anti-cancer treatment target. Cancer Immunotherapy Immune Checkpoint Immunotherapy Targeted Therapy

Reference

- Bhr C, et al. (2004) The insulin like growth factor-1 receptor (IGF-1R) as a drug target: novel approaches to cancer therapy. *Growth Horm IGF Res.* 14 (4): 287-95.
- Riedemann J, et al. (2006) IGF1R signalling and its inhibition. *Endocr Relat Cancer.* 13 Suppl 1: 33-43.
- Gualco E, et al. (2009) IGF-IR in neuroprotection and brain tumors. *Front Biosci.* 14: 352-75.
- Annenkov A. (2009) The insulin-like growth factor (IGF) receptor type 1 (IGF1R) as an essential component of the signalling network regulating neurogenesis. *Mol Neurobiol.* 40 (3): 195-215.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481