

VISTA Protein, Cynomolgus, Recombinant (His)

General Information

Synonyms:	V-set immunoregulatory receptor;C9H10orf54
Protein Construction:	Phe33-Thr192
Species:	Cynomolgus
Expression Host:	HEK293 Cells
Accession:	F7F6S6
Molecular Weight:	19 kDa (predicted); 40-60 kDa (reducing conditon due to glycosylation)

QC Testing

Biological Activity:	Immobilized Cynomolgus/Rhesus macaque B7-H5, His Tag at 1 µg/ml (100 µl/Well) on the plate. Dose response curve for Anti-B7-H5 Antibody, hFc Tag with the EC50 of 4.2 ng/ml determined by ELISA
Purity:	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

Preparation and Storage

Reconstitution:

Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

VSIR (V-Set Immunoregulatory Receptor, also known as VISTA) is a Protein Coding gene. VISTA is an immunoregulatory receptor that inhibits the T-cell response. It may promote differentiation of embryonic stem cells, by inhibiting BMP4 signaling. VSIR, or V-set immunoregulatory receptor, could be involved in the pathogenesis of chronic rhinosinusitis with nasal polyps. V-domain Immunoglobulin Suppressor of T cell Activation (VISTA) is an inhibitory immune-checkpoint molecule that suppresses CD4+ and CD8+ T cell activation

A DRUG SCREENING EXPERT

when expressed on antigen-presenting cells. VSIR is broadly expressed in the spleen, bone marrow, and other tissues. Diseases associated with VSIR include Ichthyosis, Congenital, Autosomal Recessive 6, and Monckeberg Arteriosclerosis. An important paralog of this gene is VSIG8.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481