

Fas Ligand Protein, Mouse, Recombinant (His)

General Information

Synonyms:	CD178;Apt1lg1;Fasl;Cd95l;Faslg;Tnfsf6;gld;Fas ligand;CD95-L;Fas-L
Protein Construction:	A DNA sequence encoding the mouse Faslg (AAB02915.1) (Pro132-Leu279) was expressed with a polyhistidine tag at the N-terminus. Predicted N terminal: His
Species:	Mouse
Expression Host:	P. pastoris (Yeast)
Accession:	AAB02915.1
Molecular Weight:	18.2 kDa (predicted)

QC Testing

Biological Activity:	Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 90 % as determined by SDS-PAGE.
Endotoxin:	Please contact us for more information.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:	A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.
Stability & Storage:	It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots. <small>Actual storage temperature shall be subject to the COA.</small>
Shipping:	In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Fas Ligand, also known as FASLG and CD95L, is the ligand for FAS. It is a transmembrane protein which binds to TNFRSF6/FAS. Interaction of FAS with fas Ligand is critical in triggering apoptosis of some types of cells such as lymphocytes. Fas Ligand may be involved in cytotoxic T-cell mediated apoptosis and in T-cell development. TNFRSF6/FAS-mediated apoptosis may have a role in the induction of peripheral tolerance, in the antigen-stimulated suicide of mature T-cells, or both.

Reference

Pitti R M. et al., 1998, Nature. 396 (6712): 699-703.

Hane M. et al., 1995, FEBS Lett. 373 (3): 265-8.

Schneider P. et al., 1997, J Biol Chem. 272 (30): 18827-33.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only. Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481