

PD-L1 Protein, Human, Recombinant (mFc)

General Information

Synonyms:	CD274 molecule;B7-H1;PDL1;PDCD1L1;B7H1;B7-H;PDCD1LG1;PD-L1
Protein Construction:	A DNA sequence encoding the human CD274 (NP_054862.1) (Met1-Thr239) was expressed with the Fc region of mouse IgG1 at the C-terminus. Predicted N terminal: Phe 19
Species:	Human
Expression Host:	HEK293 Cells
Accession:	Q9NZQ7-1
Molecular Weight:	51.7 kDa (predicted); 66 kDa (reducing condition, due to glycosylation)

QC Testing

Biological Activity:	<ol style="list-style-type: none">1. Measured by its binding ability in a functional ELISA. Immobilized recombinant Human PD1-His at 10 µg/ml (100 µl/well) can bind human PD-L1 with a linear range of 1.28-20 µg/ml.2. Labeled biotin to PD-L1 Protein, Human, Recombinant (ECD, Fc Tag) by a certain molar ratio;Using the Octet RED System, the affinity constant (Kd) of PD-L1 Protein, Human, Recombinant (ECD, Fc Tag), Biotinylated bound to Atezolizumab was 0.1 nM.
Purity:	> 95 % as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from 20 mM Tris, 150 mM NaCl, 10% Glycerol, pH 8.5. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:
A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Programmed death-1 ligand-1 (PD-L1, CD274, B7-H1) has been identified as the ligand for the immunoinhibitory receptor programmed death-1 (PD1/PDCD1) and has been demonstrated to play a role in the regulation of immune responses and peripheral tolerance. PD-L1/B7-H1 is a member of the growing B7 family of immune molecules and this protein contains one V-like and one C-like Ig domain within the extracellular domain, and together with PD-L2, are two ligands for PD1 which belongs to the CD28/CTLA4 family expressed on activated lymphoid cells. By binding to PD1 on activated T-cells and B-cells, PD-L1 may inhibit ongoing T-cell responses by inducing apoptosis and arresting cell-cycle progression. Accordingly, it leads to growth of immunogenic tumor growth by increasing apoptosis of antigen specific T cells and may contribute to immune evasion by cancers. PD-L1 thus is regarded as promising therapeutic target for human autoimmune disease and malignant cancers.

Cancer Immunotherapy
Co-inhibitory Immune Checkpoint Targets
Immune Checkpoint
Immune Checkpoint Blockade: Blocking Antibody
Immune Checkpoint Blockade: PD-L1 / B7-H1 / C
Immune Checkpoint Detection: Antibodies
Immune Checkpoint Detection: ELISA Antibodies
Immune Checkpoint Detection: FCM Antibodies
Immune Checkpoint Detection: ICC Antibodies
Immune Checkpoint Detection: IHC Antibodies
Immune Checkpoint Detection: WB Antibodies
Immune Checkpoint Proteins
Immune Checkpoint Targets
Immunotherapy
PD-L1 / B7-H1 / CD274
Immune Checkpoint Prote
Targeted Therapy

Reference

- Iwai Y, et al. (2002) Involvement of PD-L1 on tumor cells in the escape from host immune system and tumor immunotherapy by PD-L1 blockade. *Proc Natl Acad Sci U S A.* 99(19): 12293-7.
- Ghebeh H, et al. (2006) The B7-H1 (PD-L1) T lymphocyte-inhibitory molecule is expressed in breast cancer patients with infiltrating ductal carcinoma: correlation with important high-risk prognostic factors. *Neoplasia.* 8(3): 190-8.
- Salih HR, et al. (2006) The role of leukemia-derived B7-H1 (PD-L1) in tumor-T-cell interactions in humans. *Exp Hematol.* 34(7): 888-94.
- Wilcox RA, et al. (2009) B7-H1 (PD-L1, CD274) suppresses host immunity in T-cell lymphoproliferative disorders. *Blood.* 114(10): 2149-58.
- Ruggiero A, et al. (2009) Crystal structure of PD-L1, a ribosome inactivating protein from *Phytolacca dioica* L. leaves with the property to induce DNA cleavage. *Biopolymers.* 91(12): 1135-42.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481