

M-CSF/CSF1 Protein, Cynomolgus, Rhesus, Recombinant (His)

General Information

Synonyms:	colony stimulating factor 1 (macrophage)
Protein Construction:	A DNA sequence encoding the cynomolgus / rhesus CSF1 (XP_005542489.1) (Met1-Asn190) was expressed with a polyhistidine tag at the C-terminus. Cynomolgus and Rhesus CSF1 sequences are identical. Predicted N terminal: Glu 33
Species:	Cynomolgus,Rhesus
Expression Host:	HEK293 Cells
Accession:	XP_005542489.1
Molecular Weight:	19.8 kDa (predicted)

QC Testing

Biological Activity:	Measured in a cell proliferation assay using M-NFS-60 mouse myelogenous leukemia lymphoblast cell . The ED50 for this effect is typically 2-10 ng/mL.
Purity:	> 95 % as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:
A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Macrophage colony-stimulating factor 1, also known as CSF-1, M-CSF, Lanimostim and CSF1, is a single-pass membrane protein which is disulfide-linked as a homodimer or heterodimer. Granulocyte / macrophage colony-stimulating factors are cytokines that act in hematopoiesis by controlling the production, differentiation, and function of 2 related white cell populations of the blood, the granulocytes and the monocytes-macrophages. M-

CSF/CSF-1 is known to facilitate monocyte survival, monocyte-to-macrophage conversion, and macrophage proliferation. M-CSF/CSF-1 is a secreted cytokine which influences hemopoietic stem cells to differentiate into macrophages or other related cell types. It binds to the Colony stimulating factor 1 receptor. M-CSF/CSF-1 may also be involved in development of the placenta. The active form of M-CSF/CSF-1 is found extracellularly as a disulfide-linked homodimer, and is thought to be produced by proteolytic cleavage of membrane-bound precursors. M-CSF/CSF-1 induces cells of the monocyte/macrophage lineage. It also plays a role in immunological defenses, bone metabolism, lipoproteins clearance, fertility and pregnancy. Upregulation of M-CSF/CSF-1 in the infarcted myocardium may have an active role in healing not only through its effects on cells of monocyte/macrophage lineage, but also by regulating endothelial cell chemokine expression.

Reference

- Pandit J. et al., 1992, Science. 258: 1358-62.
Tokai M. et al., 2000, J Bacteriol. 182 (10): 2865-8.
Fan X. et al., 2001, Am J Physiol Endocrinol Metab. 280 (1): E103-11.
Frangogiannis NG. et al., 2003, Am J Physiol Heart Circ Physiol. 285 (2): H483-92.
Cupp JS. et al., 2007, Am J Surg Pathol. 31 (6): 970-6.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481