

ICOS ligand Protein, Cynomolgus, Recombinant (hFc)

General Information

Synonyms:	inducible T-cell co-stimulator ligand
Protein Construction:	A DNA sequence encoding the cynomolgus ICOSLG (XP_005548617.1) (Met1-Thr256) was expressed with the Fc region of human IgG1 at the C-terminus. Predicted N terminal: Asp 19
Species:	Cynomolgus
Expression Host:	HEK293 Cells
Accession:	XP_005548617.1
Molecular Weight:	53.3 kDa (predicted)

QC Testing

Biological Activity:	Immobilized Recombinant Cynomolgus ICOS Ligand / B7-H2 Protein (Fc Tag) at 2 µg/ml (100 µl/well) can bind Recombinant Human ICOS / AILIM / CD278 Protein (His & Fc Tag), the EC50 is 8-36 ng/mL.
Purity:	> 95 % as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Inducible co-stimulator ligand (ICOSL), also known as B7-H2, is a member of the B7 family of co-stimulatory molecules related to B7-1 and B7-2. It is a transmembrane glycoprotein with extracellular IgV and IgC domains and binds to ICOS on activated T cells, thus delivers a positive costimulatory signal for optimal T cell function. The structural features of ICOSL are crucial for its costimulatory function. The present study shows that ICOSL displays a

marked oligomerization potential, resembling more like B7-1 than B7-2. B7-H2-dependent signaling may play an active role in a proliferative response rather than in cytokine and chemokine production. The CD28/B7 and ICOS/B7-H2 pathways are both critical for costimulating T cell immune responses. Deficiency in either pathway results in defective T cell activation, cytokine production, and germinal center formation. Cancer Immunotherapy Co-stimulatory Immune Checkpoint Targets Immune Checkpoint Immune Checkpoint Detection: Antibodies Immune Checkpoint Detection: ELISA Antibodies Immune Checkpoint Detection: FCM Antibodies Immune Checkpoint Detection: WB Antibodies Immune Checkpoint Targets Immunotherapy Targeted Therapy

Reference

- Flesch IE. (2002) Inducible costimulator-ligand (ICOS-L). *J Biol Regul Homeost Agents*. 16(3): 217-9.
- Kajiwara K, et al. (2009) Expression and function of the inducible costimulator ligand B7-H2 in human airway smooth muscle cells. *Allergol Int*. 58(4): 573-83.
- Wong SC, et al. (2009) Functional hierarchy and relative contribution of the CD28/B7 and ICOS/B7-H2 costimulatory pathways to T cell-mediated delayed-type hypersensitivity. *Cell Immunol*. 256(1-2): 64-71.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481