

PDGFB Protein, Mouse, Recombinant (His)

General Information

Synonyms:	platelet-derived growth factor β polypeptide;Sis;platelet-derived growth factor beta polypeptide;PDGF-B
Protein Construction:	A DNA sequence encoding the mouse PDGFB (AAH23427.1) (Ser82-Thr190) was expressed with a polyhistidine tag at the N-terminus. Predicted N terminal: His
Species:	Mouse
Expression Host:	P. pastoris (Yeast)
Accession:	Q8R3X9
Molecular Weight:	14.2 kDa (predicted)

QC Testing

Biological Activity:	Measured by its binding ability in a functional ELISA. Immobilized Human PDGFRB at 2 μ g/ml (100 μ l/well) can bind Mouse PDGF-B His, the EC50 of Mouse PDGF-B His is 4-24 ng/mL.
Purity:	> 95 % as determined by SDS-PAGE.
Endotoxin:	Please contact us for more information.
Formulation:	Lyophilized from a solution filtered through a 0.22 μ m filter, containing 30% CAN, 0.1% TFA. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Platelet-derived growth factor-B (PDGFB) is necessary for normal cardiovascular development. The administration of PDGFB alone normalized tumor vasculature by increasing periendothelial coverage and vascular functionality. Interestingly, this effect exerted by PDGFB was also observed in the presence of DAPT. So PDGFB is able to improve tumor vascularity and allows the anticancer action of DAPT in the tumor.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481