

## TIGIT Protein, Cynomolgus, Recombinant (hFc)

### General Information

Synonyms:	T cell immunoreceptor with Ig and ITIM domains
Protein Construction:	A DNA sequence encoding the cynomolgus TIGIT (XP_015300911.1) (Met1-Pro209) was expressed with the Fc region of human IgG1 at the C-terminus. Predicted N terminal: Lys 27
Species:	Cynomolgus
Expression Host:	HEK293 Cells
Accession:	A0A7N9I9V4
Molecular Weight:	47.2 kDa (predicted)

### QC Testing

Biological Activity:	Immobilized human CD155/PVR Protein, Human, Recombinant (ECD, His Tag) at 2µg/mL (100µL/well) can bind TIGIT Protein, Cynomolgus, Recombinant (ECD, hFc Tag), the EC50 of TIGIT Protein, Cynomolgus, Recombinant (ECD, hFc Tag)(90890-C02H) is 5-30 ng/mL.
Purity:	> 95 % as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

### Preparation and Storage

**Reconstitution:**  
Reconstituted with sterile deionized water to 0.25 mg/mL. Reconstitution conditions may vary depending on the lot.

#### Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

#### Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

### Protein Background

TIGIT, also known as V-set and transmembrane domain-containing protein 3 (VSTM3) or V-set and immunoglobulin domain-containing protein 9 (VSIG9) is a new surface protein containing an immunoglobulin variable domain, a transmembrane domain and an immunoreceptor tyrosine-based inhibitory motif (ITIM). TIGIT is expressed on regulatory, memory, activated T cells and NK cells. It binds PVR with high affinity, and PVRL2 with

lower affinity, but not PVRL3. Knockdown of TIGIT with siRNA in human memory T cells did not affect T cell responses, however, TIGIT inhibits NK cytotoxicity directly through its ITIM. TIGIT suppresses T cell activation by promoting the generation of mature immunoregulatory dendritic cells. The binding of PVR to TIGIT on human dendritic cells enhanced the production of IL-1 and diminished the production of IL-12p4. Also, TIGIT counter inhibits the NK-mediated killing of tumor cells and protects normal cells from NK-mediated cytotoxicity thus providing an "alternative self" mechanism for MHC class I inhibition. Cancer Immunotherapy Co-inhibitory Immune Checkpoint Targets Immune Checkpoint Immune Checkpoint Targets Immunotherapy Targeted Therapy

### Reference

Ota T., et al., (2004), Complete sequencing and characterization of 21,243 full-length human cDNAs. Nat. Genet. 36: 40-45.

Bechtel S., et al., (2007), The full-ORF clone resource of the German cDNA consortium. BMC Genomics 8:399-399.

Yu X., Harden K., et al., (2009), The surface protein TIGIT suppresses T cell activation by promoting the generation of mature immunoregulatory dendritic cells. Nat. Immunol. 10:48-57.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E\_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481