

## VSIG4 Protein, Human, Recombinant (hFc), Biotinylated

### General Information

Synonyms:	CRIg;Z39IG;V-set and immunoglobulin domain containing 4
Protein Construction:	A DNA sequence encoding the human VSIG4 (NP_009199.1) (Met1-Pro283) was expressed with the Fc region of human IgG1 at the C-terminus. The purified protein was biotinylated in vitro. Predicted N terminal: Arg 20
Species:	Human
Expression Host:	HEK293 Cells
Accession:	Q9Y279-1
Molecular Weight:	56.2 kDa (predicted)

### QC Testing

Biological Activity:	Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 90 % as determined by SDS-PAGE
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

### Preparation and Storage

**Reconstitution:**  
A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

**Stability & Storage:**

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

**Shipping:**

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

### Protein Background

VSIG4 (V-set and immunoglobulin domain containing 4), also known as complement receptor of the immunoglobulin superfamily (CRIg) and Z39Ig, is a type I transmembrane glycoprotein. It is a B7 family-related protein and an Ig superfamily member. In contrast to the B7 family members which contain two IgG domains, VSIG4 contains one complete V-type I g domain and a truncated C-type I g domain. VSIG4 is exclusively expressed

## A DRUG SCREENING EXPERT

---

on tissue resident macrophages and binds to multimers of C3b and iC3b that are covalently attached to particle surfaces. No VSIG4 expression appears to be present in T and B cells. VSIG4 functions as a negative regulator of T cell activation, and may be involved in the maintenance of peripheral T cell tolerance, and is also identified as a potent suppressor of established inflammation. Mouse VSIG4 is synthesized as a 28 amino acid precursor that contains a signal sequence, a V-type I g domain (aa 36-115), one potential N-linked glycosylation site, and a single transmembrane domain. The V-type I g domain of mouse VSIG4 shares 86% and 8% aa sequence identity with the V-type I g domains of rat and human VSIG4, respectively.

### Reference

- Vogt, L. et al., 2006, J Clin Invest. 116: 2817-2826.  
Helmy, K. et al., 2006, Cell. 124:915-927.  
Wiesmann, C. et al., 2006, Nature. 444:217-220.  
Zang, X. et al., 2006, J Clin Invest. 116: 2590-2593.  
Katschke, KJ. et al., 2007, J. Exp. Med. 204:1319-1325.  
He, JQ. et al., 2008, Mol Immunol. 45: 4041-4047.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E\_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481