

Human Papilloma Virus type 11 (HPV 11) L1 Protein-VLP

General Information

Species:	HPV
Expression Host:	Baculovirus Insect Cells
Molecular Weight:	55 kDa (predicted)

QC Testing

Biological Activity:	EM Images: Negative staining electron microscopy of Human Papilloma Virus type 11 L1 virus-like particles (VLPs)(Routinely tested).
Purity:	≥ 90 % as determined by SDS-PAGE.
Endotoxin:	< 0.01 EU/μg of the protein as determined by the LAL method.
Formulation:	Supplied as sterile 50 mM His, 0.5M NaCl, pH 6. 2.

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store the product under sterile conditions at -70°C or lower. Samples are stable for up to 12 months at -80°C. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

Proteins are shipped with blue ice.

Protein Background

Papillomaviruses are highly species-specific and can cause squamous epithelial and fibroepithelial tumors in their hosts. Human papillomaviruses (HPVs) are associated with benign and malignant hyperproliferation of cells, with a wide variety of clinical manifestations ranging from condyloma acuminatum to cervical carcinoma. HPV infection is the most common sexually transmitted disease. More than 4 HPV types so far identified are known to infect the genital tract. Genital HPVs are divided into 'low risk' HPVs such as HPV 6 and 11 and 'high risk' HPV types such as 16, 18, 31, 33, 35, 39, 45 and 52, 58 which are responsible for more than 95% of HPV-induced cervical cancer. Vaccination against these high-risk types seems to be the most feasible prevention for cervical cancer. Indeed, clinical trials have shown prophylactic HPV vaccines to be effective against HPV infection, cervical intraepithelial neoplasia (CIN), and genital warts, but protection is type-specific and the currently developed vaccines target only a few types. These vaccines are based on papillomavirus-like particles (VLPs) composed of the major capsid protein, L1. The L1 protein self assembles into VLPs when expressed at high levels in eukaryotic or insect cells. VLPs are composed of 36 copies of L1 protein organized into 72 pentamers, so-called capsomeres, to form particles that are immunologically indistinguishable from native virions. Experimentally induced VLP antisera are mostly type-specific for neutralization. Minor cross-neutralization has been observed only between closely related

HPV types, e.g. HPV6 and 11, HPV18 and 45, or HPV16 and 31. Structure analysis has revealed the presence of several hypervariable loops on the outer surface of the capsid. With a few exceptions, all HPV-neutralizing monoclonal antibodies analyzed so far are type-specific and recognize conformational epitopes within surface-exposed hypervariable loops of the major capsid protein L1.

Reference

- Zur Hausen, H., 1989, Cancer Res. 49: 4677-4691.
Chan, SY, et al., 1995, J. Virol. 69: 3074-3083.
Lorincz AT, et al., 1992, Obstet Gynecol. 79: 328-337.
Munoz N, et al., 2003, N Engl J Med. 348: 518-527.
Pastrana DV, et al., 2004, Virology. 321: 205-216.
Christensen ND, et al., 1996, Virology. 224: 477-486.
Combata AL, et al., 2002, J Virol. 76: 6480-6486.
Christensen ND, et al., 2001, Virology. 291:324-334.
Fleury MJ, et al., Arch Virol. 2006, 151: 1511-1523.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481