

CSF1R Protein, Mouse, Recombinant (hFc)

General Information

Synonyms:	AI323359;Csfmr;Fim-2;Fms;colony stimulating factor 1 receptor;M-CSF-R;M-CSFR;CD115;CSF-1R
Protein Construction:	A DNA sequence encoding the mouse CSF1R (NP_001032948.2) (Met1-Ser511) was expressed with the Fc region of human IgG1 at the C-terminus. Predicted N terminal: Ala 20
Species:	Mouse
Expression Host:	HEK293 Cells
Accession:	P09581
Molecular Weight:	81.9 kDa (predicted)

QC Testing

Biological Activity:	Measured by its ability to inhibit the mouse CSF-induced proliferation of M-NFS-60 mouse myelogenous leukemia lymphoblast cells. The ED50 for this effect is typically 30-140ng/mL in the presence of 15 ng/mL mouse M-CSF.
Purity:	> 95 % as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

M-CSFR encoded by the proto-oncogene *c-fms* is the receptor for colony stimulating factor 1 (CSF1R), a cytokine involved in the proliferation, differentiation, and activation of macrophages. This cell surface glycoprotein is consisted by an extracellular ligand-binding domain, a single membrane-spanning segment, and an intracellular

tyrosine kinase domain. Binding of CSF1 activates the receptor kinase, leading to "autophosphorylation" of receptor subunits and the concomitant phosphorylation of a series of cellular proteins on tyrosine residues. CSF1R is a tyrosine kinase receptor that is absolutely required for macrophage differentiation and thus occupies a central role in hematopoiesis. CSF1 and its receptor (CSF1R, product of c-fms proto-oncogene) were initially implicated as essential for normal monocyte development as well as for trophoblastic implantation. This apparent role for CSF1/CSF1R in normal mammary gland development is very intriguing because this receptor/ligand pair has also been found to be important in the biology of breast cancer in which abnormal expression of CSF1 and its receptor correlates with tumor cell invasiveness and adverse clinical prognosis. Tumor cell expression of CSF1R is under the control of several steroid hormones (glucocorticoids and progestins) and the binding of several bHLH transcription factors, while tumor cell expression of CSF-1 appears to be regulated by other hormones, some of which are involved in normal lactogenic differentiation. However, studies have demonstrated that CSF1 and CSF1R have additional roles in mammary gland development during pregnancy and lactation. The role of CSF1 and CSF1R in normal and neoplastic mammary development that may elucidate potential relationships of growth factor-induced biological changes in the breast during pregnancy and tumor progression.

Reference

- Sherr CJ. (1990) The colony-stimulating factor 1 receptor: pleiotropy of signal-response coupling. *Lymphokine Res.* 9(4): 543-8.
- Kacinski BM. (1997) CSF-1 and its receptor in breast carcinomas and neoplasms of the female reproductive tract. *Mol Reprod Dev.* 46(1): 71-4.
- Sapi E, et al. (1999) The role of CSF-1 in normal and neoplastic breast physiology. *Proc Soc Exp Biol Med.* 220(1): 1-8.
- Sapi E. (2004) The role of CSF-1 in normal physiology of mammary gland and breast cancer: an update. *Exp Biol Med (Maywood).* 229(1): 1-11.
- Bonifer C, et al. (2008) The transcriptional regulation of the Colony-Stimulating Factor 1 Receptor (*csf1r*) gene during hematopoiesis. *Front Biosci.* 13: 549-60.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481