

## Transferrin Receptor/TFRC Protein, Human, Recombinant (His), Biotinylated

### General Information

Synonyms:	TRFR;p90;TR;TFR;CD71;TFR1;T9;transferrin receptor
Protein Construction:	A DNA sequence encoding the human TFRC (NP_003225.2) (Cys 89-Phe 760) was expressed with a polyhistidine tag at the N-terminus. The purified protein was biotinylated in vitro. Predicted N terminal: His
Species:	Human
Expression Host:	HEK293 Cells
Accession:	P02786
Molecular Weight:	77.4 kDa (predicted)

### QC Testing

Biological Activity:	Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 85 % as determined by SDS-PAGE
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

### Preparation and Storage

**Reconstitution:**  
A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

**Stability & Storage:**

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

**Shipping:**

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

### Protein Background

Transferrin receptor protein 1, also known as transferrin receptor, Trfr, p9, CD71 and TFRC, is a single-pass type II membrane protein that belongs to the peptidase M28 family and M28B subfamily. TFRC / CD71 is a membrane-bound protein expressed in larger amounts in proliferating. The specific expression of TFRC can represent a diagnostic tool or a therapeutic target in solid tumours expressing this antigen. Transferrin receptor is necessary

## A DRUG SCREENING EXPERT

---

for development of erythrocytes and the nervous system. TFRC / CD71 is regulated by cellular iron levels through binding of the iron regulatory proteins, IRP1 and IRP2, to iron-responsive elements in the 3'-UTR. Up-regulated upon mitogenic stimulation. TFRC / CD71 represents a marker of malignant transformation in the pancreas that could be applied as potential diagnostic and therapeutic target.

### Reference

Douabin-Gicquel V., et al., 2001, Hum. Genet. 109:393-401.

Ryschich, E. et al., 2004, Eur J Cancer. 40 (9):1418-22.

Tosoni D., et al., 2005, Cell 123:875-888.

Wollscheid B., et al., 2009, Nat. Biotechnol. 27:378-386.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E\_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481