

## LAG-3 Protein, Mouse, Recombinant

### General Information

Synonyms:	lymphocyte-activation gene 3;LAG-3;Ly66;CD223
Protein Construction:	A DNA sequence encoding the mouse LAG3 (NP_032505.1) (Met1-Leu442) was expressed. Predicted N terminal: Gly 24
Species:	Mouse
Expression Host:	HEK293 Cells
Accession:	Q61790
Molecular Weight:	45.25 kDa (predicted)

### QC Testing

Biological Activity:	Measured by its ability to bind FGL1 Protein, Mouse, Recombinant (hFc Tag) in functional ELISA.
Purity:	> 95 % as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

### Preparation and Storage

#### Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

#### Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

#### Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

### Protein Background

LAG3 (Lymphocyte Activating 3) is a Protein Coding gene. 2 alternatively spliced human isoforms have been reported. LAG3, also known as CD223 and Lymphocyte activation gene 3, belongs to the immunoglobulin (Ig) superfamily. The LAG3 gene contains 8 exons. It is selectively expressed in activated T and NK cells. LAG3 contains 4 extracellular Ig-like domains and has a negative regulatory function in T cells. It also acts as a new marker of T cell-induced B cell activation. As a soluble molecule, LAG3 activates antigen-presenting cells through MHC class II

signaling, leading to increased antigen-specific T-cell responses in vivo. Diseases associated with LAG3 include Smoldering Myeloma and Kyphoscoliotic Heart Disease. Cancer Immunotherapy Co-inhibitory Immune Checkpoint Targets Immune Checkpoint Immune Checkpoint Targets Immunotherapy Targeted Therapy

### Reference

Sigrid Hannier. et al., 1998, The Journal of Immunology. 161(8): 4058-65.

Triebel F. et al., 1990, J Exp Med. 171 (5): 1393-405.

Baixeras E. et al., 1992, J Exp Med. 176 (2): 327-37.

Huard B. et al., 1997, Proc Natl Acad Sci. 94 (11): 5744-9.

Kisielow M. et al., 2005, Eur J Immunol. 35 (7): 2081-8.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E\_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481