

Podoplanin Protein, Human, Recombinant (His)

General Information

Synonyms:	T1A-2;PA2.26;GP36;T1A;AGGRUS;OTS8;T1A2;TI1A;HT1A-1;GP40;Gp38;podoplanin
Protein Construction:	A DNA sequence encoding the human PDPN (Q86YL7) (Met1-Lys123) was expressed with a polyhistidine tag at the C-terminus. Predicted N terminal: Ala 23
Species:	Human
Expression Host:	HEK293 Cells
Accession:	Q86YL7
Molecular Weight:	11.78 kDa (predicted)

QC Testing

Biological Activity:	Measured by its ability to bind recombinant CLEC1B-His/Biotin in a functional ELISA.
Purity:	> 95 % as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:
A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Podoplanin, also known as PDPN, is a type-I integral membrane glycoprotein with diverse distribution in human tissues. The physiological function of this protein may be related to its mucin-type character. The homologous protein in other species has been described as a differentiation antigen and influenza-virus receptor. The specific function of this protein has not been determined. Alternatively spliced transcript variants encoding different isoforms have been identified. PDPN is a mucin-type glycoprotein negatively charged by extensive O-glycosylation

and a high content of sialic acid, which expresses the adhesive property. It is selectively expressed in lymphatic endothelium as well as lymphangiomas, Kaposi sarcomas, and in a subset of angiosarcomas with probable lymphatic differentiation. PDPN may contribute to form odontoblastic fiber or function as the anchorage to the tooth development and in proliferating epithelial cells of cervical loop and apical bud. The intensity of podoplanin expression is negatively correlated with the expression of CD34 and factor VIII. Podoplanin would be useful as a diagnostic marker for epithelioid hemangioendothelioma in liver tumors. Cancer Immunotherapy Immune Checkpoint Immunotherapy Targeted Therapy

Reference

- Kimura, N. et al., 2005, Pathol Int. 55 (2): 83-86.
Ordóñez, N.G., 2006, Adv Anat Pathol. 13 (2): 83-88.
Wicki, A. et al., 2007, Br. J. Cancer. 96 (1): 1-5.
Fujii, T. et al., 2008, Mod Pathol. 21 (2): 125-130.
Sawa, Y. et al., 2008, Acta Histochem Cytochem. 41 (5): 121-126.
Kaddu, S. et al., 2009, Am J Dermatopathol. 31 (2): 137-139.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481