

## Influenza A H3N2 (A/Kansas/14/2017) Neuraminidase/NA Protein (His)

### General Information

Synonyms:	NA Protein
Protein Construction:	A DNA sequence encoding the Influenza A virus (A/Kansas/14/2017 (H3N2)) neuraminidase (AVG71505.1) (Ile30-Ile469) was expressed with a vasodilator-stimulated phosphoprotein tetramerization domain at the N-terminus and a polyhistidine tag at the C-terminus. Predicted N terminal: Asp
Species:	H3N2
Expression Host:	Baculovirus Insect Cells
Accession:	A0A2L2FIU4
Molecular Weight:	54.6 kDa (predicted); 56.5 kDa (reducing conditions)

### QC Testing

Biological Activity:	1.Measured by its ability to cleave a fluorogenic substrate, 2'-(4-Methylumbelliferyl)- $\alpha$ -D-N-acetylneuraminic acid. 2.Immobilized Recombinant H3N2 (A/Kansas/14/2017) NA Protein (His Tag) (Cat#TMPY-06066) at 1 $\mu$ g/mL (100 $\mu$ L/well) can bind Anti-Influenza NA (Clone 1G01), the EC50 is 6-18 ng/mL (QC tested).
Purity:	> 90 % as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/ $\mu$ g of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 $\mu$ m filter, containing 20 mM Tris, 300 mM NaCl, pH 8.5, 10% glycerol. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

### Preparation and Storage

Reconstitution:	Reconstituted with sterile deionized water to 0.15 mg/mL. Reconstitution conditions may vary depending on the lot.
Stability & Storage:	It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots. <small>Actual storage temperature shall be subject to the COA.</small>
Shipping:	In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

### Protein Background

Neuraminidases are enzymes that cleave sialic acid groups from glycoproteins. Influenza neuraminidase is a type of neuraminidase found on the surface of influenza viruses that enables the virus to be released from the host cell. Influenza neuraminidase is composed of four identical subunits arranged in a square. It is normally attached to the virus surface through a long protein stalk. The active sites are in a deep depression on the upper surface. They bind to polysaccharide chains and clip off the sugars at the end. The surface of neuraminidase is decorated with several polysaccharide chains that are similar to the polysaccharide chains that decorate our cell surface proteins. Neuraminidase (NA) and hemagglutinin (HA) are major membrane glycoproteins found on the surface of the influenza virus. Hemagglutinin binds to the sialic acid-containing receptors on the surface of host cells during initial infection and at the end of an infectious cycle. Neuraminidase, on the other hand, cleaves the HA-sialic acid bondage from the newly formed virions and the host cell receptors during budding. Neuraminidase thus is described as a receptor-destroying enzyme that facilitates virus release and efficient spread of the progeny virus from cell to cell. Influenza antibody and influenza antibodies are very important research tools for influenza diagnosis, influenza vaccine development, and anti-influenza virus therapy development. The monoclonal or polyclonal antibody can be raised with protein based antigen or peptide-based antigen. Antibodies raised with protein-based antigen could have better specificity and/or binding affinity than antibodies raised with peptide based antigen, but the cost associated with the recombinant protein antigen is usually higher. Anti-influenza virus hemagglutinin (HA) monoclonal antibody or polyclonal antibody can be used for ELISA assay, western blotting detection, Immunohistochemistry (IHC), flow cytometry, neutralization assay, hemagglutinin inhibition assay, and early diagnosis of influenza viral infection. Sino Biological has developed state-of-the-art monoclonal antibody development technology platforms: mouse monoclonal antibody and rabbit monoclonal antibody. Our rabbit monoclonal antibody platform is one of a kind and offers some unique advantages over mouse monoclonal antibodies, such as high affinity, low cross-reactivity with rabbit polyclonal antibodies.

### Reference

- Sardet C., et al., (1989), Molecular cloning, primary structure, and expression of the human growth factor-activatable Na<sup>+</sup>/H<sup>+</sup> antiporter. *Cell* 56:271-280.
- Sardet C., et al., (1990), Growth factors induce phosphorylation of the Na<sup>+</sup>/H<sup>+</sup> antiporter, glycoprotein of 110 kD. *Science* 247:723-726.
- Tse C.-M., et al., (1991), Molecular cloning and expression of a cDNA encoding the rabbit ileal villus cell basolateral membrane Na<sup>+</sup>/H<sup>+</sup> exchanger. *EMBO J.* 10:1957-1967.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

**This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use**

Tel: 781-999-4286    E\_mail: info@targetmol.com    Address: 34 Washington Street, Wellesley Hills, MA 02481