

CD155/PVR Protein, Human, Recombinant (hFc & Avi), Biotinylated

General Information

Synonyms:	TAGE4;CD155;NECL5;poliovirus receptor;HVED;Necl-5;PVS
Protein Construction:	Trp21-Asn343
Species:	Human
Expression Host:	HEK293 Cells
Accession:	P15151-1
Molecular Weight:	62.82 kDa (Predicted); 80-100 kDa (Reducing conditions due to glycosylation)

QC Testing

Biological Activity:	Immobilized Biotinylated Human CD155, hFc Tag at 5µg/ml (100µl/Well) on streptavidin (5µg/ml) precoated plate. Dose response curve for Human TIGIT, His Tag with the EC50 of 1.14µg/ml determined by ELISA (QC Test).
Purity:	> 95% as determined by Bis-Tris PAGE
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 µg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

CD155 is a cell surface adhesion molecule functioning in tumor cell migration, invasion, and metastasis, and not surprisingly, is also designated as a common tumor-associated antigen. CD155 is also recognized by NK cells to induce their cytotoxicity. CD155 is also commonly referred to as the "poliovirus receptor," or PVR.

Reference

- Freistadt MS, et al. (2000) Hematopoietic cells from CD155-transgenic mice express CD155 and support poliovirus replication ex vivo. *Microb Pathog.* 29(4): 203-12.
- Sato T, et al. (2004) Involvement of heterophilic trans-interaction of Necl-5/Tage4/PVR/CD155 with nectin-3 in formation of nectin- and cadherin-based adherens junctions. *Genes Cells.* 9(9): 791-9.
- Kakunaga S, et al. (2004) Enhancement of serum- and platelet-derived growth factor-induced cell proliferation by Necl-5/Tage4/poliovirus receptor/CD155 through the Ras-Raf-MEK-ERK signaling. *J Biol Chem.* 279(35): 36419-25.
- Sato T, et al. (2005) Common signaling pathway is used by the trans-interaction of Necl-5/Tage4/PVR/CD155 and nectin, and of nectin and nectin during the formation of cell-cell adhesion. *Cancer Sci.* 96(9): 578-89.
- Minami Y, et al. (2007) Involvement of up-regulated Necl-5/Tage4/PVR/CD155 in the loss of contact inhibition in transformed NIH3T3 cells. *Biochem Biophys Res Commun.* 352(4): 856-60.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481