

CD48 Protein, Human, Recombinant (His & Avi), Biotinylated

General Information

Synonyms:	SLAMF2;BLAST;mCD48;hCD48;Blast-1;MEM-102;BLAST1;BCM1;CD48 molecule
Protein Construction:	Gln27-Ser220
Species:	Human
Expression Host:	HEK293 Cells
Accession:	P09326-1
Molecular Weight:	25.25 kDa (predicted); 45-60 kDa (reducing conditions, due to glycosylation)

QC Testing

Biological Activity:	Immobilized Biotinylated Human CD48, His-Avi Tag at 2 µg/ml (100 µl/well) on the streptavidin precoated plate (5 µg/ml). Dose response curve for Human 2B4, hFc Tag with the EC50 of 0.9 ng/ml determined by ELISA. (QC Test)
Purity:	> 95% as determined by Bis-Tris PAGE; > 95% as determined by HPLC
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 µg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

CD48, also known as BLAST-1, BCM-1, and SLAMF2, is a 65 kDa GPI-linked protein in the CD2 family of immunoglobulin superfamily molecules. CD2 and 2B4 (CD244) are known ligands for CD48. CD48 protein is expressed on most lineage-committed hematopoietic cells but not on hematopoietic stem cells or multipotent hematopoietic progenitors.

Reference

Messmer B, et al. (2006) CD48 stimulation by 2B4 (CD244)-expressing targets activates human NK cells. J Immunol. 176(8): 4646-50.

Milstein O, et al. (2008) Nanoscale increases in CD2-CD48-mediated intermembrane spacing decrease adhesion and reorganize the immunological synapse. J Biol Chem. 283(49): 34414-22.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481