

## CEACAM6 Protein, Human, Recombinant (His & Avi), Biotinylated

### General Information

Synonyms:	NCA;CEAL;CD66c;carcinoembryonic antigen-related cell adhesion molecule 6 (non-specific cross reacting antigen);CEA
Protein Construction:	A DNA sequence encoding the Human CEACAM6 (NP_002474.3) (Met1-Gly320) was expressed with a C-terminal polyhistidine tag followed by an AVI tag. The expressed protein was biotinylated in vivo by the Biotin-Protein ligase (BirA enzyme) which is co-expressed. Predicted N terminal: Lys 35
Species:	Human
Expression Host:	HEK293 Cells
Accession:	NP_002474.3
Molecular Weight:	34.47 kDa (predicted); 79.81 kDa (reducing conditions)

### QC Testing

Biological Activity:	Immobilized TPP-3310 derived Anti-CEACAM-6 antibody at 2 µg/mL (100 µL/well) can bind Recombinant Human CEACAM6 Protein (His & Avi Tag), Biotinylated (Cat#TMPY-06758), the EC50 is 1.7-5.1 ng/mL (QC tested).
Purity:	≥ 95 % as determined by SDS-PAGE. ≥ 95 % as determined by SEC-HPLC.
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

### Preparation and Storage

Reconstitution:	A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.
Stability & Storage:	It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots. <small>Actual storage temperature shall be subject to the COA.</small>
Shipping:	In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

### Protein Background

Carcinoembryonic antigen-related cell adhesion molecule 6 (CEACAM6), also known as nonspecific crossreacting

antigen (NCA) and CD66c, is one of seven human CEACAM family members within the immunoglobulin superfamily. It is a glycosylphosphatidylinositol-linked immunoglobulin superfamily member that is overexpressed in a variety of human cancers, including colon, breast and lung and is associated with tumorigenesis, tumour cell adhesion, invasion and metastasis. CEACAM6 is a unique mediator of migration and invasion of drug resistant oestrogen-deprived breast cancer cells, and this protein could be an important biomarker of metastasis. CEACAM6 is expressed by granulocytes and their progenitors. It is also expressed by epithelia of various organs and is upregulated in pancreatic and colon adenocarcinomas, as well as hyperplastic polyps. Resistance to adhesion-related apoptosis in tumor cells is conferred in the condition of CEACAM6 overexpression.

### Reference

Duxbury MS, et al. (2004) Overexpression of CEACAM6 promotes insulin-like growth factor I-induced pancreatic adenocarcinoma cellular invasiveness. *Oncogene*. 23(34): 5834-42.

Lewis-Wambi JS, et al. (2008) Overexpression of CEACAM6 promotes migration and invasion of oestrogen-deprived breast cancer cells. *Eur J Cancer*. 44(12): 1770-9.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286    E\_mail: info@targetmol.com    Address: 34 Washington Street, Wellesley Hills, MA 02481