

IGFBP-7 Protein, Mouse, Recombinant (aa 26-281, His)

General Information

Synonyms:	Mac25;insulin-like growth factor binding protein 7;AGM;Fstl2
Protein Construction:	A DNA sequence encoding the Human IGFBP7 (AAH92538.1) (Ser26-Leu281) was expressed, with a polyhistidine tag at the N-terminus. Predicted N terminal: His
Species:	Mouse
Expression Host:	HEK293 Cells
Accession:	AAH92538.1
Molecular Weight:	27.69 kDa (predicted); 33.36 kDa and 24.38 kDa (reducing conditions)

QC Testing

Biological Activity:	Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	≥ 90 % as determined by SDS-PAGE. ≥ 95 % as determined by SEC-HPLC.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing 25 mM MES, 150 mM NaCl, 0.02% Tween 20, 5% Trehalose, 10% Glycerol, pH 5. 5. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:
A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Insulin-like growth factor-binding protein 7 (IGFBP7) is a member of the IGFBP family. It has been identified in colorectal adenocarcinoma (CRC) cell lines. The Insulin-like growth factor-binding protein also known as IGFBP serves as a carrier protein for Insulin-like growth factor 1. IGFBPs are distinct but are sharing regions with strong homology. All members of the IGFBP family bind IGF-I and IGF-II with about equal affinity. Insulin-like growth

factor (IGF) binding proteins (IGFBPs) have been shown to either inhibit or enhance the action of IGF or act in an IGF-independent manner in the prostate. IGFBP7 could inhibit cell growth, decrease soft agar colony formation activity, and induce apoptosis in RKO and SW620 cells. There is mounting evidence that the structure of the IGFBP proteins plays a key role in the regulation of IGF bioavailability, by modulating its molecular size, capillary membrane permeability, target tissue specificity, cell membrane adherence, and IGF affinity.

Reference

- Oh Y, et al. (1996) Synthesis and characterization of insulin-like growth factor-binding protein (IGFBP)-7. Recombinant human mac25 protein specifically binds IGF-I and -II. *J Biol Chem.* 271(48): 30322-5.
- Wilson EM, et al. (1997) Generation and characterization of an IGFBP-7 antibody: identification of 31kD IGFBP-7 in human biological fluids and Hs578T human breast cancer conditioned media. *J Clin Endocrinol Metab.* 82(4): 1301-3.
- Lin J, et al. (2007) Methylation patterns of IGFBP7 in colon cancer cell lines are associated with levels of gene expression. *J Pathol.* 212(1): 83-90.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481