


NU 7026

Chemical Properties

CAS No. : 154447-35-5
 Formula: C17H15NO3
 Molecular Weight: 281.31
 Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year
Actual storage temperature shall be subject to the COA.



Biological Description

Description	NU 7026 (DNA-PK Inhibitor II) is an effective DNA-PK inhibitor (IC50: 0.23 μ M, in cell-free assays), 60-fold selective for DNA-PK than PI3K and no inhibition against both ATM and ATR.
Targets(IC50)	Apoptosis,ATM/ATR,DNA-PK,PI3K
In vitro	The bioavailability of NU7026, administered intraperitoneally (i.p.) or orally (p.o.) at a dose of 20 mg/kg, was found to be 20% and 15%, respectively. In mice, NU7026 administered intravenously (i.v.) at the same dosage exhibited a rapid plasma clearance rate of 0.108/h.
In vivo	NU7026 inhibits DNA DSB repair in the V3YAC cell line by 56%. In the CLL cell line (I83) and primary CLL lymphocytes, NU7026 exhibits synergistic cytotoxic activity with Bendamustine at concentrations below 10 μ M. In K562 cells, NU7026 (10 μ M) enhances the growth inhibitory effects of doxorubicin, amsacrine, idarubicin, aminobenzotriazole, etoposide, and mitoxantrone, with PF50 values ranging approximately from 19 (for mAMSA) to 2 (for idarubicin). Also, at 10 μ M, NU7026 enhances the growth inhibition induced by etoposide in leukemia cells (PF50: 10.53) and the etoposide-induced G2 phase cell cycle arrest in K562 cells. A 4-hour exposure to NU7026 (10 μ M) combined with 3 Gy radiation is necessary for a significant radiosensitization effect in CH1 human ovarian cancer cells. In the I83 cell line, NU7026 (10 μ M) increases G(2)/M phase arrest and γ H2AX throughout the cell cycle, both induced by Bendamustine, as well as apoptosis. Finally, at 55 μ M, NU7026 notably induces telomere fusion in p53-deficient MEFs and causes less telomere fusion in MEFs lacking both p53 and Ligase IV.
Kinase Assay	Mammalian DNA-PK (500 ng/ μ L) is isolated from HeLa cell nuclear extract after chromatography using Q-Sepharose, S-Sepharose, and Heparin agarose. DNA-PK (250 ng) activity is measured at 30°C, in a final volume of 40 μ L, in buffer containing 25 mM HEPES (pH 7.4), 12.5 mM MgCl ₂ , 50 mM KCl, 1 mM DTT, 10% v/v Glycerol, 0.1% w/v NP-40, and 1 mg of the substrate GST-p53N66 (the NH ₂ -terminal 66 amino acid residues of human wild-type p53 fused to glutathione S-transferase) in polypropylene 96-well plates. To the assay mix, varying concentrations of inhibitor (in DMSO at a final concentration of 1% v/v) are added. After 10 min of incubation, ATP is added to give a final concentration of 50 μ M, along with a 30-mer double-stranded DNA oligonucleotide (final concentration of 0.5 ng/mL), to initiate the reaction. After 1 h with shaking, 150 μ L of PBS are added to the reaction, and 5 μ L are then transferred to a 96-well opaque white plate containing 45 μ L of PBS per well, where the GSTp53N66 substrate is allowed

Kinase Assay	to bind to the wells for 1 h. To detect the phosphorylation event on the serine 15 residue of p53 elicited by DNA-PK, a p53 phosphoserine-15 antibody is used in a basic ELISA procedure. An antirabbit horseradish peroxidase-conjugated secondary antibody is then used in the ELISA before the addition of chemiluminescence reagent to detect the signal as measured by chemiluminescent counting via a TopCount NXT[1].
Cell Research	NU7026 is dissolved in DMSO and stored, and then diluted with appropriate media before use[2]. I83 cells are plated in RPMI 1640 medium with 10% FBS (1.5×10 ⁵ cells/mL) and treated with vehicle (DMSO), 5 μM CLB, CLB IC50, 10 μM NU7026, or the combination of both drugs for 0, 6, 24, and 48 h. Cell cycle distribution, apoptosis, DNA-PK phosphorylation, and γH2AX determination are determined, and they are expressed as a percentage of cells in each phase of the cycle. DNA content is analyzed with a FACSCalibur flow cytometer equipped with CellQuest software[2].

Solubility Information

Solubility	DMSO: 2.69 mg/mL (9.56 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	--

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	3.5548 mL	17.774 mL	35.548 mL
5 mM	0.711 mL	3.5548 mL	7.1096 mL
10 mM	0.3555 mL	1.7774 mL	3.5548 mL
50 mM	0.0711 mL	0.3555 mL	0.711 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

- Veuger SJ, et al. Cancer Res, 2003, 63(18), 62008-62015.
- Willmore E, et al. Blood, 2004, 103(12), 4659-4665.
- Nutley BP, et al. Br J Cancer, 2005, 93(9), 12011-12018.
- Amrein L, et al. J Pharmacol Exp Ther, 2007, 321(3), 848-855.
- Williams ES, et al. Cancer Res, 2009, 69(5), 2100-2107.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481