


Cisapride

Chemical Properties

| | |
|-------------------|---|
| CAS No. : | 81098-60-4 |
| Formula: | C ₂₃ H ₂₉ ClFN ₃ O ₄ |
| Molecular Weight: | 465.95 |
| Storage: | Powder: -20°C for 3 years In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA. |



Biological Description

| | |
|---------------|---|
| Description | Cisapride (R 51619) is a serotonin type 4 (5-HT ₄) receptor agonists are potent prokinetic agents that act on serotonin receptors in the intestine and promote intestinal peristalsis, increase gastric emptying and decrease esophageal reflux. |
| Targets(IC50) | 5-HT Receptor, Potassium Channel |
| In vitro | Cisapride inhibits vascular Kv current independently of serotonin 5-HT ₄ -receptor activation[2]. As HERG channel blocker, cisapride, can inhibit the growth of gastric cancer cells by altering distribution of cell cycle and inducing apoptosis so as to be of potential value in the treatment of gastric cancer. Cisapride could inhibit the growth and clonogenicity of human gastric cancer lines by specific blockage of HERG channel in a time- and dose-dependent manner while has little effects on GES cells[3]. |
| Cell Research | Cells in the logarithmic growth phase are harvested and seeded in 96-well plates. The cell number is diluted to 5000/well in 200 microliters of RPMI1640 medium. In vitro experiments are carried out on the gastric cancer cell lines and GES cell line, using an MTT proliferation assay. MTT stains live cells, and the optical density absorbance at 490 nm wavelength correlates with the cell number. Cisapride is added to the cells at concentrations of 0, 12.5, 25, 50, 100 or 200 nM and equivalent volume of ethanol is added as control in order to eliminate the effects of solvent. Twenty microliters of MTT is added to each well of the cell culture 4 h before termination of the culture at 37°C. One hundred and fifty microliters of dimethyl sulfoxide is added to each well at the end of the culture and the plate is agitated for 10 minutes. The absorbency at 490 nm is read by a BIOHIT BP800 plate reader. Growth curve is drawn according to MTT colorimetry. The inhibition rate is calculated.(Only for Reference) |

Solubility Information

| | |
|---------------------|---|
| Solubility | DMSO: 255 mg/mL (547.27 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble) |
| In vivo Formulation | 10% DMSO+40% PEG300+5% Tween 80+45% Saline: 2 mg/mL (4.29 mM),Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may</i> |

A DRUG SCREENING EXPERT

| | |
|---------------------|---|
| In vivo Formulation | <i>vary and should be modified based on specific experimental conditions.</i> |
|---------------------|---|

Preparing Stock Solutions

| | 1mg | 5mg | 10mg |
|-------|-----------|------------|------------|
| 1 mM | 2.1462 mL | 10.7308 mL | 21.4615 mL |
| 5 mM | 0.4292 mL | 2.1462 mL | 4.2923 mL |
| 10 mM | 0.2146 mL | 1.0731 mL | 2.1462 mL |
| 50 mM | 0.0429 mL | 0.2146 mL | 0.4292 mL |

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Park JS, et al. Eur J Med Chem. 2016, 109:75-88.

Kim HW, et al. Biochem Biophys Res Commun. 2016, 478(3):1423-8.

Shao XD, et al. Cancer Biol Ther. 2005, 4(3):295-301.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481