


Piceatannol 3'-O-glucoside

Chemical Properties

CAS No. :	94356-26-0
Formula:	C ₂₀ H ₂₂ O ₉
Molecular Weight:	406.38
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



Biological Description

Description	Piceatannol 3'-O-glucoside (Piceatannol 3-glycoside) increases pulmonary blood oxygen tension and decreases pulmonary interstitial edema, so it can protect lung injury.
Targets(IC50)	Arginase,NO Synthase

Solubility Information

Solubility	DMSO: 60 mg/mL (147.65 mM),Sonication is recommended. Pyridine, Methanol, etc.: Soluble, Ethanol: Soluble, (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 2 mg/mL (4.92 mM),Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.4608 mL	12.3038 mL	24.6075 mL
5 mM	0.4922 mL	2.4608 mL	4.9215 mL
10 mM	0.2461 mL	1.2304 mL	2.4608 mL
50 mM	0.0492 mL	0.2461 mL	0.4922 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Woo A, et al. Arginase inhibition by piceatannol-3'-O- β -D-glucopyranoside improves endothelial dysfunction via activation of endothelial nitric oxide synthase in ApoE-null mice fed a high-cholesterol diet. *Int J Mol Med*. 2013 Apr;31(4):803-10.

Woo A, et al. Piceatannol-3'-O-beta-D-glucopyranoside as an active component of rhubarb activates endothelial nitric oxide synthase through inhibition of arginase activity. *Exp Mol Med*. 2010 Jul 31;42(7):524-32.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481