



## Solubility Information

Solubility	DMSO: 115 mg/mL (181.81 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 3.3 mg/mL (5.22 mM),Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

## Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	1.5809 mL	7.9046 mL	15.8093 mL
5 mM	0.3162 mL	1.5809 mL	3.1619 mL
10 mM	0.1581 mL	0.7905 mL	1.5809 mL
50 mM	0.0316 mL	0.1581 mL	0.3162 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

## Reference

Tavazoie MF, et al. LXR/ApoE Activation Restricts Innate Immune Suppression in Cancer. Cell. 2018 Feb 8;172(4):825-840.e18.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481