

Bithionol sulfoxide

Chemical Properties


CAS No. : 844-26-8

Formula: C₁₂H₆Cl₄O₃S

Molecular Weight: 372.05

Storage: Store at low temperature, Keep away from moisture
Powder: -20°C for 3 years | In solvent: -80°C for 1 year

Actual storage temperature shall be subject to the COA.



Biological Description

Description	Bithionol sulfoxide (Bithionoloxide) is a compound with inhibitory effects on parasites, suppressing the activity of <i>Schistosoma japonicum</i> and <i>Fasciola hepatica</i> .
Targets(IC50)	Parasite
In vitro	Bithionol sulfoxide (0.1-10 mg/mL, 72 hours) showed toxicity to <i>Neoparamoeba</i> parasites in seawater. [1]
In vivo	Bithionol sulfoxide significantly inhibited C.S. administration The oviposition of a worm. Bithionol sulfoxide showed a slight insecticidal effect at high doses of 100 mg/kg. [2] Bithionol sulfoxide is mutagenic against <i>Salmonella typhimurium</i> strains TA98 and TA100. [3]

Solubility Information

Solubility	DMSO: 200 mg/mL (537.56 mM), Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 5 mg/mL (13.44 mM), Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.6878 mL	13.4391 mL	26.8781 mL
5 mM	0.5376 mL	2.6878 mL	5.3756 mL
10 mM	0.2688 mL	1.3439 mL	2.6878 mL
50 mM	0.0538 mL	0.2688 mL	0.5376 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Florent RL, et al. In vitro toxicity of bithionol and bithionol sulphoxide to *Neoparamoeba* spp., the causative agent of amoebic gill disease (AGD). *Dis Aquat Organ*. 2010 Sep 17;91(3):257-62.

Kang SY, et al. [A Study On The Chemotherapy In Clonorchiasis: Report 1. An Experimental Study On Chemotherapy With Dithiazanine Iodide And Bithionol Sulfoxide In Clonorchiasis]. *Kisaengchunghak Chapchi*. 1965 Jun;3(1):19-30. Korean.

Mourot D, Mourot A. Mutagenicity of bithionol sulfoxide and its metabolites in the salmonella/mammalian microsome test. *Mutat Res*. 1987 May;188(1):53-5.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481