


Dp44mT

## Chemical Properties

CAS No. : 152095-12-0  
 Formula: C<sub>14</sub>H<sub>15</sub>N<sub>5</sub>S  
 Molecular Weight: 285.37  
 Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year  
*Actual storage temperature shall be subject to the COA.*



## Biological Description

Description	Dp44mT, a effective iron chelator, has selective antitumor activity.
Targets(IC50)	Apoptosis,Ferroptosis
In vitro	In MDA-MB-231 cells, Dp44mT selectively targets towards topoisomerase topo2 $\alpha$ resulting in DNA damage. Dp44mT inhibits antiproliferative effects significantly with IC50 of 30 nM, 60 nM, and 60 nM for SK-N-MC, SK-Mel-28, and MCF-7 cells, respectively, while showing no effects on normal MRC-5 fibroblasts. In SK-N-MC neuroepithelioma and M109 cells,Dp44mT inhibits cellular Fe uptake from Fe-Tf , and induces cell apoptosis. Dp44mT, as a Pgp substrate, also overcomes multidrug resistance by the hijacking of lysosomal P-glycoprotein (Pgp).
In vivo	Dp44mT (0.4 mg/kg, i.v.) inhibits M109 tumor implanted in CD2F1 mice by a dose-dependent manner.
Cell Research	SK-N-MC, SK-Mel-28, MCF-7 and MRC-5 cells are incubated in the presence and absence of DFO, 311, 3-aminopyridine-2-carboxaldehyde thiosemicarbazone, doxorubicin, and the DpT series of chelators (0-25 $\mu$ M) for 72 hours at 37°C. The effect of the chelators on proliferation is examined using the MTT assay.
Animal Research	CD2F1 mice that are implanted M109 tumors intravenous inject Dp44mT (dissolved in Propylene glycol) 4 mg/kg in twice daily.

## Solubility Information

Solubility	DMSO: 16.67 mg/mL (58.42 mM),Sonication is recommended. Ethanol: 47 mg/mL (164.7 mM),Sonication is recommended. H2O: Insoluble, (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 2 mg/mL (7.01 mM),Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

### Preparing Stock Solutions

---

	1mg	5mg	10mg
1 mM	3.5042 mL	17.5211 mL	35.0422 mL
5 mM	0.7008 mL	3.5042 mL	7.0084 mL
10 mM	0.3504 mL	1.7521 mL	3.5042 mL
50 mM	0.0701 mL	0.3504 mL	0.7008 mL

---

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

### Reference

Yuan J, et al. Blood. 2004, 104(5), 1450-1458.

Rao VA, et al. Cancer Res. 2009, 69(3), 948-957.

Jansson PJ, et al. J Biol Chem. 2015, 290(15), 9588-9603.

Whitnall M, et al. A class of iron chelators with a wide spectrum of potent antitumor activity that overcomes resistance to chemotherapeutics. Proc Natl Acad Sci U S A. 2006 Oct 3;103(40):14901-6.

Yalowich J C , Wu X , Zhang R , et al. The anticancer thiosemicarbazones Dp44mT and triapine lack inhibitory effects as catalytic inhibitors or poisons of DNA topoisomerase II $\alpha$ [J]. Biochemical Pharmacology, 2012, 84(1):52-58.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481