


Latifoline

Chemical Properties

CAS No. :	4438-06-6
Formula:	C ₂₂ H ₃₅ NO
Molecular Weight:	329.52
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



Biological Description

Description	Latifoline is a alkaloid from Funtumia latifolia Stapf. barks.
Targets(IC50)	Others

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	3.0347 mL	15.1736 mL	30.3472 mL
5 mM	0.6069 mL	3.0347 mL	6.0694 mL
10 mM	0.3035 mL	1.5174 mL	3.0347 mL
50 mM	0.0607 mL	0.3035 mL	0.6069 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

- Drutu I, Krygowski ES, Wood JL. Reactive enols in synthesis 2. Synthesis of (+)-latifolic acid and (+)-latifoline. J Org Chem. 2001 Oct 19;66(21):7025-9. PubMed PMID: 11597224.
- Hincks JR, Kim HY, Segall HJ, Molyneux RJ, Stermitz FR, Coulombe RA Jr. DNA cross-linking in mammalian cells by pyrrolizidine alkaloids: structure-activity relationships. Toxicol Appl Pharmacol. 1991 Oct;111(1):90-8. PubMed PMID: 1949039.
- JANOT MM, QUI KH, GOUTAREL R. [Steroid alkaloids. Structure of latifoline, a new alkaloid from Funtumia latifolia Stapf. barks]. C R Hebd Seances Acad Sci. 1962 Feb 12;254:1326-8. French. PubMed PMID: 14451176.
- Kim HY, Stermitz FR, Molyneux RJ, Wilson DW, Taylor D, Coulombe RA Jr. Structural influences on pyrrolizidine alkaloid-induced cytopathology. Toxicol Appl Pharmacol. 1993 Sep;122(1):61-9. PubMed PMID: 8378933.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481