


Omipalisib

Chemical Properties

| | |
|-------------------|--|
| CAS No. : | 1086062-66-9 |
| Formula: | C ₂₅ H ₁₇ F ₂ N ₅ O ₃ S |
| Molecular Weight: | 505.5 |
| Storage: | Store at low temperature Powder: -20°C for 3 years In solvent: -80°C for 1 year <small>Actual storage temperature shall be subject to the COA.</small> |



Biological Description

| | |
|---------------|--|
| Description | Omipalisib (GSK2126458) is a small-molecule pyridylsulfonamide inhibitor of phosphatidylinositol 3-kinase (PI3K) with potential antineoplastic activity. |
| Targets(IC50) | Autophagy,mTOR,PI3K |
| In vitro | GSK2126458 reduces the levels of pAKT-S473 in T47D cells (IC50=0.41 nM) and BT474 cells (IC50=0.18 nM), leading to cell cycle arrest at the G1 phase and inhibiting cell proliferation. |
| In vivo | In the BT474 human xenograft model, GSK2126458 (300 µg/kg) effectively inhibits tumor cell growth in a dose-dependent manner. Clinical trials have demonstrated that GSK2126458 possesses oral bioactivity in mice, rats, dogs, and monkeys, along with favorable blood clearance properties. |
| Kinase Assay | HTRF In vitro Profiling Assays for PI3K Inhibition : Compounds are serially diluted (3-fold in 100% DMSO) across a 384-well polypropylene mother plate from column 1 to column 12 and column 13 to column 24, to yield 11 concentrations for GSK2126458. Columns 6 and 18 contain only DMSO. Once titrations are made, 0.05µL is transferred to a 384-well low-volume assay plate. This assay plate contains three pharmacological controls (known PI3K inhibitors) and 3 assay controls: (1) Enzyme without inhibitor; (2) Buffer minus enzyme, and (3) Buffer minus enzyme plus native PIP3. DMSO is stamped into all wells of columns 6 and 18. PIP3 is added at 40 µM in 1X Reaction buffer (1µL of 200 µM PIP3) to alternating rows of column 18 (wells 18 B, D, F, H, J, L, N, P). The no-enzyme control reactions are run in wells 18 A, C, E, G, I, K, M, O (0.1µL of 100% DMSO). The PI3-Kinase profiling assay is optimized using the HTRF kit. The assay kit contains seven reagents: 1) 4X Reaction Buffer; 2) native PIP2 (substrate); 3) Stop A (EDTA); 4) Stop B (Biotin-PIP3); 5) Detection Mix A (Streptavidin-APC); 6) Detection Mix B (Eu-labeled Anti-GST plus GST-tagged PHdomain); 7) Detection Mix C (KF). PI3Kinase Reaction Buffer is prepared by diluting the stock 1:4 with de-ionized water. Freshly prepared DTT is added at a final concentration of 5 mM on the day of use. Enzyme addition and compound pre-incubation are initiated by the addition of 2.5µL of PI3K (at twice its final concentration) in 1X reaction buffer to all wells using a Multidrop Combi. Plates are incubated at room temperature for 15 minutes. Reactions are initiated by addition of 2.5µL of 2X substrate solution (PIP2 and ATP in 1X reaction buffer) using a Multidrop Combi. Plates are incubated at room temperature for one hour. Reactions are quenched by the addition of |

| | |
|---------------|--|
| Kinase Assay | 2.5µL of stop solution (Stop A and Stop B pre-mixed at a ratio of 5:1, respectively) to all wells using the Multidrop Combi. The quenched reactions are then processed to detect product formation by adding 2.5µL of Detection Solution to all wells using the Multidrop Combi (Detection mix C, Detection mix A, and Detection mix B combined together in an 18:1:1 ratio, i.e.: for a 6000 µL total volume, mix 5400 µL Detection mix C, 300µL Detection mix A, and 300 µL Detection mix B. Note: this solution should be prepared 2 hours prior to use). Following a one hour incubation in the dark, the HTRF signal is measured on the Envision plate reader set for 330 nM excitation and dual emission detection at 620 nM (Eu) and 665 nM (APC). |
| Cell Research | BT474, HCC1954 and T-47D (human breast) are cultured in RPMI-1640 containing 10% fetal bovine serum at 37 °C in 5% CO2 incubator. Cells are split into T75 flask two to three days prior to assay set up at density which yields approximately 70-80% confluence at time of harvest for assay. Cells are harvested using 0.25% trypsin-EDTA. Cell counts are performed on cell suspension using Trypan Blue exclusion staining. Cells are then plated in 384 well black flat bottom polystyrene in 48 µL of culture media per well at 1,000 cells/well. All plates are placed at 5% CO2, 37 °C overnight and GSK2126458 is added the following day. One plate is treated with CellTiter-Glo for a day 0 (t=0) measurement and read as described below. GSK2126458 is prepared in clear bottom polypropylene 384 well plates with consecutive two fold dilutions. 4 µL of these dilutions are added to 105 µL culture media, after mixing the solution, 2 µL of these dilutions are added into each well of the cell plates. The final concentration of DMSO in all wells is 0.15%. Cells are incubated at 37 °C, 5% CO2 for 72 hours. Following 72 hours of incubation with GSK2126458 each plate is developed and read. CellTiter-Glo reagent is added to assay plates using a volume equivalent to the cell culture volume in the wells. Plates are shaken for approximately two minutes and incubated at room temperature for approximately 30 minutes and chemiluminescent signal is read on the Analyst GT reader. Results are expressed as a percent of the t=0 and plotted against the GSK2126458 concentration. Cell growth inhibition is determined for GSK2126458 by fitting the dose response with a 4 or 6 parameter curve fit using XLfit software and determining the concentration that inhibits 50% of the cell growth (gIC50) with the Y min as the t=0 and Y max as the DMSO control. Value from wells with no cells is subtracted from all samples for background correction.(Only for Reference) |

Solubility Information

| | |
|---------------------|---|
| Solubility | DMSO: 50 mg/mL (98.91 mM), Sonication is recommended. Ethanol: < 1 mg/mL (insoluble or slightly soluble), H2O: < 1 mg/mL (insoluble or slightly soluble) (< 1 mg/ml refers to the product slightly soluble or insoluble) |
| In vivo Formulation | 10% DMSO+40% PEG300+5% Tween 80+45% Saline: 2 mg/mL (3.96 mM), Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i> |

Preparing Stock Solutions

| | 1mg | 5mg | 10mg |
|-------|------------|------------|-------------|
| 1 mM | 1.9782 mL | 9.8912 mL | 19.7824 mL |
| 5 mM | 0.3956 mL | 1.9782 mL | 3.9565 mL |
| 10 mM | 0.1978 mL | 0.9891 mL | 1.9782 mL |
| 50 mM | 0.0396 mL | 0.1978 mL | 0.3956 mL |

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

- Knight SD, et al. ACS Med. Chem. Lett. 2010, 1 (1), 39-43.
Greger JG, et al. Mol Cancer Ther. 2012, 11(4), 909-920.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481