

Colcemid

Chemical Properties


CAS No. : 477-30-5

Formula: C₂₁H₂₅N₅O

Molecular Weight: 371.43

Storage: Keep away from moisture, Store at low temperature
Powder: -20°C for 3 years | In solvent: -80°C for 1 year

Actual storage temperature shall be subject to the COA.



Biological Description

| | |
|----------------------------|--|
| Description | Colcemid (NSC-3096) is a microtubule polymerization inhibitor, with the main target being Tubulin and an IC ₅₀ of 2.4 μM. Colcemid can induce apoptosis and can be used in the research of tumors and embryo cloning. |
| Targets(IC ₅₀) | Apoptosis, Microtubule Associated |

Solubility Information

| | |
|---------------------|---|
| Solubility | DMSO: 55 mg/mL (148.08 mM), Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble) |
| In vivo Formulation | 10% DMSO+40% PEG300+5% Tween 80+45% Saline: 2 mg/mL (5.38 mM), Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i> |

Preparing Stock Solutions

| | 1mg | 5mg | 10mg |
|-------|-----------|------------|-----------|
| 1 mM | 2.6923 mL | 13.4615 mL | 26.923 mL |
| 5 mM | 0.5385 mL | 2.6923 mL | 5.3846 mL |
| 10 mM | 0.2692 mL | 1.3461 mL | 2.6923 mL |
| 50 mM | 0.0538 mL | 0.2692 mL | 0.5385 mL |

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Gao Y, Ren J, Zhang L, Zhang Y, Wu X, Jiang H, Xu F, Yuan B, Yu X, Zhang J. The effects of demecolcine, alone or in combination with sucrose on bovine oocyte protrusion rate, MAPK1 protein level and c-mos gene expression level. *Cell Physiol Biochem*. 2014;34(6):1974-82. doi: 10.1159/000366393. Epub 2014 Nov 25.

Yin Y, Mei M, Zhang D, Zhang S, Fan A, Zhou H, Li Z. The construction of cloned Sika deer embryos (*Cervus nippon hortulorum*) by demecolcine auxiliary enucleation. *Reprod Domest Anim*. 2014 Feb;49(1):164-9. doi: 10.1111/rda.12246. Epub 2013 Oct 21.

Meng Q, Wu X, Bunch TD, White K, Sessions BR, Davies CJ, Rickords L, Li GP. Enucleation of demecolcine-treated bovine oocytes in cytochalasin-free medium: mechanism investigation and practical improvement. *Cell Reprogram*. 2011 Oct;13(5):411-8. doi: 10.1089/cell.2011.0012. Epub 2011 Jul 8.

Saraiva NZ, Perecin F, Méo SC, Ferreira CR, Tetzner TA, Garcia JM. Demecolcine effects on microtubule kinetics and on chemically assisted enucleation of bovine oocytes. *Cloning Stem Cells*. 2009 Mar;11(1):141-52. doi: 10.1089/clo.2008.0044.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481