


Quinalizarin

Chemical Properties

CAS No. :	81-61-8
Formula:	C ₁₄ H ₈ O ₆
Molecular Weight:	272.21
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



Biological Description

Description	Quinalizarin, the most selective CK2 inhibitor, is superior to CX-4945 which is the first-in-class CK2 inhibitor.
Targets(IC50)	Others, Casein Kinase

Solubility Information

Solubility	DMSO: Soluble, (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	---

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	3.6736 mL	18.3682 mL	36.7363 mL
5 mM	0.7347 mL	3.6736 mL	7.3473 mL
10 mM	0.3674 mL	1.8368 mL	3.6736 mL
50 mM	0.0735 mL	0.3674 mL	0.7347 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Meng LQ, Wang Y, Luo YH, Piao XJ, Liu C, Wang Y, Zhang Y, Wang JR, Wang H, Xu WT, Liu Y, Wu YQ, Sun HN, Han YH, Jin MH, Shen GN, Fang NZ, Jin CH. Quinalizarin Induces Apoptosis through Reactive Oxygen Species (ROS)-Mediated Mitogen-Activated Protein Kinase (MAPK) and Signal Transducer and Activator of Transcription 3 (STAT3) Signaling Pathways in Colorectal Cancer Cells. *Med Sci Monit.* 2018 Jun 3;24:3710-3719. doi: 10.12659/MSM.907163. PubMed PMID: 29860266; PubMed Central PMCID: PMC6014151.

Meng LQ, Liu C, Luo YH, Piao XJ, Wang Y, Zhang Y, Wang JR, Wang H, Xu WT, Liu Y, Wu YQ, Sun HN, Han YH, Jin MH, Shen GN, Zang YQ, Li J, Fang NZ, Cui YD, Jin CH. Quinalizarin exerts an anti-tumour effect on lung cancer A549 cells by modulating the Akt, MAPK, STAT3 and p53 signalling pathways. *Mol Med Rep.* 2018 Feb;17(2):2626-2634. doi: 10.3892/mmr.2017.8110. Epub 2017 Nov 20. PubMed PMID: 29207064.

Li Q, Li K, Yang T, Zhang S, Zhou Y, Li Z, Xiong J, Zhou F, Zhou X, Liu L, Meng R, Wu G. Association of protein kinase CK2 inhibition with cellular radiosensitivity of non-small cell lung cancer. *Sci Rep.* 2017 Nov 23;7(1):16134. doi: 10.1038/s41598-017-16012-1. PubMed PMID: 29170453; PubMed Central PMCID: PMC5700935.

Willibald M, Bayer G, Stahlhut V, Poschmann G, Stühler K, Gierke B, Pawlak M, Seeger H, Mueck AO, Niederacher D, Fehm T, Neubauer H. Progesterone receptor membrane component 1 is phosphorylated upon progestin treatment in breast cancer cells. *Oncotarget.* 2017 Aug 2;8(42):72480-72493. doi: 10.18632/oncotarget.19819. eCollection 2017 Sep 22. PubMed PMID: 29069804; PubMed Central PMCID: PMC5641147.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481