


DASA-58

Chemical Properties

CAS No. : 1203494-49-8
 Formula: C₁₉H₂₃N₃O₆S₂
 Molecular Weight: 453.53
 Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year
 Actual storage temperature shall be subject to the COA.



Biological Description

Description	DASA-58 is a specific and potent Pyruvate kinase M2 (PKM2) activator.
Targets(IC50)	PKM
In vitro	DASA-58 inhibits LPS-induced Hif-1 α and IL-1 β , as well as the expression of a range of other Hif-1 α -dependent genes in primary BMDMs, and also inhibits glycolysis and the accumulation of succinate in LPS-activated macrophages. [1] In PC3 cells, DASA-58 impairs stromal-induced EMT program by restoring PK activity and abrogating the nuclear translocation of PKM2, as well as its association with HIF-1 α . DASA-58 also dramatically reduces (~6-fold) CAFs-induced lung metastases formation in PC3 cells. [2]
In vivo	DASA-58 (40 μ M) affects EMT of prostate cancers and tumor dissemination in SCID mice. [2]
Kinase Assay	CDK kinase assay: Kinase assays are performed in 96-well polypropylene plates. Each reaction contained 2 μ g of histone H1 at a final concentration of 10 μ M [γ -33P]ATP (0.2 μ Ci/well; approximately twice the experimentally determined Km), 10 mM MgCl ₂ , 1 mM DTT, 0.01% Triton X-100, and 10% glycerol in a 40 μ L volume. The reaction is initiated with the addition of 20 μ L enzyme (6 ng cdk2/well resulting in a final concentration of 1.6 nM), which is previously diluted 1:50–1:200 in the same buffer, and allowed to proceed for 1 h at room temperature. Reaction is stopped by the addition of 0.01 mL 10% phosphoric acid, and 25 μ L of reaction mixture is transferred to P30 phosphocellulose filter mat paper. The filter mat is washed three times with 1.0% phosphoric acid, air dried, and then counted for radioactivity in a liquid scintillation counter. The cdk4 kinase assay for cyclin D1-cdk4 is carried out in a polypropylene 96-well microtiter plate format measuring the incorporation of radioactive phosphate into GST-Rb. Purified cyclin D1-cdk4 is incubated with 1 μ g GST-Rb in 20 mM HEPES (pH 7.5) in the presence of 10 mM MgCl ₂ , 1 mM DTT, 0.01% Triton X-100, and 10% glycerol. The final cdk4 concentration is 10 ng/well, or 1.6 nM. Kinase reaction is initiated by the addition of ATP at a final concentration of 10 μ M ATP (twice the experimentally determined Km) and [γ -33P]ATP (1.0 μ Ci per well) in a 60- μ L volume and allowed to proceed at room temperature for 1 h. Reaction is stopped by the addition of 0.01 ml 10% phosphoric acid, and 25 μ L of reaction mixture is transferred to P30 phosphocellulose filter mat paper. The filter mat is treated as for Cdk1/Cdk2 assays.

Solubility Information

Solubility	Ethanol: < 1 mg/mL (insoluble or slightly soluble), DMSO: 125.5 mg/mL (276.72 mM),Sonication is recommended. H2O: < 1 mg/mL (insoluble or slightly soluble), (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 3.3 mg/mL (7.28 mM),Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.2049 mL	11.0246 mL	22.0493 mL
5 mM	0.441 mL	2.2049 mL	4.4099 mL
10 mM	0.2205 mL	1.1025 mL	2.2049 mL
50 mM	0.0441 mL	0.2205 mL	0.441 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Palsson-McDermott EM, et al. Cell Metab. 2015, 21(1), 65-80.

Yang L, Zhang J, Hu C, et al.Nuclear translocation of PKM2 mediates keratinocyte metabolic reprogramming in psoriasis.Experimental Dermatology.2023

Giannoni E, et al. Oncotarget. 2015, 6(27), 24061-24074.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481