


## Sarcosine

## Chemical Properties

CAS No. :	107-97-1
Formula:	C <sub>3</sub> H <sub>7</sub> NO <sub>2</sub>
Molecular Weight:	89.09
Storage:	Powder: -20°C for 3 years   In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



## Biological Description

Description	Sarcosine (Methylglycine) is a competitive inhibitor of the type I glycine transporter (GlyT1) and an N-methyl-D-aspartate receptor (NMDAR) co-agonist.
Targets(IC50)	Endogenous Metabolite, GlyT
In vitro	Sarcosine is a GlyR agonist in addition to being a GlyT1 inhibitor and NMDAR co-agonist, but it is less potent than glycine as a GlyR agonist and is not a full agonist[1]. The viability of the sarcosine-treated cells is significantly reduced[4].
In vivo	Sarcosine has weak anticonvulsant properties[2]. It ameliorates (prepulse inhibition)PPI deficits in mGluR5 knockout mice[3].
Cell Research	Immediately after the cells grow to 50-60% confluence, the cultivation medium is replaced by fresh medium to synchronise cell growth. Cells are cultivated for 24 h under these conditions. Subsequently, the culture medium is supplemented with sarcosine (N-methylglycine) diluted to a final concentration 10, 150, 250, 500, 1,000 and 1,500 µM. Treatment is carried out for 0, 6, 12, 24 and 72 h, and samples are collected at these strictly defined time points.(Only for Reference)

## Solubility Information

Solubility	DMSO: Insoluble, H <sub>2</sub> O: 250 mg/mL (2806.15 mM), Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	---

### Preparing Stock Solutions

---

	<b>1mg</b>	<b>5mg</b>	<b>10mg</b>
1 mM	11.2246 mL	56.123 mL	112.246 mL
5 mM	2.2449 mL	11.2246 mL	22.4492 mL
10 mM	1.1225 mL	5.6123 mL	11.2246 mL
50 mM	0.2245 mL	1.1225 mL	2.2449 mL

---

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

### Reference

Zhang HX, et al. *Neuropharmacology*. 57(5-6):551-5.

Socała K, et al. *Pharmacol Rep*. 2010, 62(2):392-7.

Chen HH, et al. *Psychopharmacology (Berl)*. 2010, 209(4):343-50.

Sudhakaran P R , Binu S , Soumya S J . Effect of sarcosine on endothelial function relevant to angiogenesis.[J]. *Journal of Cancer Research & Therapeutics*, 2014, 10(3):603-610.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286

E\_mail:info@targetmol.com

Address:34 Washington Street,Wellesley Hills,MA 02481