


Galeterone

Chemical Properties

CAS No. :	851983-85-2
Formula:	C ₂₆ H ₃₂ N ₂ O
Molecular Weight:	388.55
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



Biological Description

Description	Galeterone (VN 124) is an orally bioavailable small-molecule androgen receptor modulator and CYP17 lyase inhibitor with potential antiandrogen activity. Galeterone exhibits three distinct mechanisms of action: 1) as an androgen receptor antagonist, 2) as a CYP17 lyase inhibitor and 3) by decreasing overall androgen receptor levels in prostate cancer tumors, all of which may result in a decrease in androgen-dependent growth signaling. Localized to the endoplasmic reticulum (ER), the cytochrome P450 enzyme CYP17 (P450C17 or CYP17A1) exhibits both 17 α -hydroxylase and 17, 20-lyase activities, and plays a key role in the steroidogenic pathway that produces progestins, mineralocorticoids, glucocorticoids, androgens, and estrogens.
Targets(IC50)	Androgen Receptor, Cytochromes P450
In vitro	Galeterone is effective at preventing binding of [3H]-R1881 to the mutant LNCaP AR (T877A) with IC ₅₀ of 845 nM. Galeterone inhibits the DHT-induced proliferation of LNCaP and LAPC4 cells in a dose-dependent manner with IC ₅₀ of 6 μ M and 3.2 μ M, respectively. [1] Galeterone also inhibits the binding of [3H]-R1881 to the T575A mutant AR in PC3 cells with IC ₅₀ of 454 nM. Galeterone potently inhibits the proliferation of LNCaP and LAPC4 cells in the absence of DHT stimulation with IC ₅₀ of 2.6 μ M and 4 μ M, respectively. Furthermore, Galeterone treatment increases the degradation rate of the AR in a dose-dependent manner. [2] Galeterone potently inhibits the growth of the androgen-independent cell lines PC-3 and DU-145 in a dose-dependent manner with GI ₅₀ of 7.82 μ M and 7.55 μ M, respectively. Galeterone induces the endoplasmic reticulum stress response resulting in down-regulation of cyclin D1 protein expression and cyclin E2 mRNA. [3] Galeterone effectively inhibits proliferation of HP-LNCaP and C4-2B cell lines with IC ₅₀ of 2.9 μ M and 9.7 μ M, respectively. Galeterone treatment at 1 μ M effectively inhibits androgen receptor activation in LNCaP cells (50%) and HP-LNCaP cells (70%). Galeterone decreases activation of the androgen receptor in both LNCaP cells and HP-LNCaP cells with IC ₅₀ of 1 μ M and 411 nM, respectively, and down-regulates androgen receptor protein expression by 50% after 24 hour of treatment. [4] Galeterone reduces AR protein and mRNA expression, antagonizes AR-dependent promoter activation induced by androgen, and significantly reduces the phospho-4EBP1 levels. [6]
In vivo	Administration of Galeterone at 50 mg/kg twice daily is very effective at inhibiting the growth of androgen-dependent LAPC4 human prostate tumor xenograft, with a 93.8% reduction in the mean final tumor volume compared with controls, and it is also significantly more effective than castration. [1] Treatment of Galeterone (0.13 mM/kg

In vivo	twice daily) or Galeterone (0.13 mmol/kg twice daily) plus castration induces regression of LAPC4 tumor xenografts in SCID mice by 26.55% and 60.67%, respectively. Treatments with Galeterone or Galeterone plus castration causes marked reduction in AR protein of 10- and 5-fold, respectively. [2]
Kinase Assay	In vitro assay of CYP17: The in vitro CYP17 inhibitory activity of Galeterone is evaluated using rapid acetic acid releasing assay (AARA), utilizing intact P450c17-expressing E. coli as the enzyme source. It involves the use of [21-3H]-17 α -hydroxypregnenolone as the substrate, and CYP17 activity is measured by the amount of tritiated acetic acid formed during the cleavage of the C-21 side chain of the substrate. IC50 value is obtained directly from plots relating percentage inhibition versus inhibitor concentration over appropriate ranges.
Cell Research	Cells are seeded in 24 well multi-well plates. Cells are treated with the increasing concentration of Galeterone in steroid free medium with or without 1 nM DHT (LNCaP), or 10 nM DHT (LAPC4) and allowed to grow for 7 days. The number of viable cells is compared by MTT assay (LAPC4) or XTT assay (LNCaP) on the 7th day. (Only for Reference)

Solubility Information

Solubility	Ethanol: 19.4 mg/mL (49.93 mM),Sonication is recommended. DMSO: 19.4 mg/mL (49.93 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 2 mg/mL (5.15 mM),Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.5737 mL	12.8684 mL	25.7367 mL
5 mM	0.5147 mL	2.5737 mL	5.1473 mL
10 mM	0.2574 mL	1.2868 mL	2.5737 mL
50 mM	0.0515 mL	0.2574 mL	0.5147 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Handratta VD, et al. J Med Chem, 2005, 48(8), 2972-2984.

Xiao M, Ha S, Zhu J, et al. Structure-Activity Relationship (SAR) Studies of Novel Monovalent AR/AR-V7 Dual Degraders with Potent Efficacy against Advanced Prostate Cancer. Journal of Medicinal Chemistry. 2024

Vasaitis T, et al. Mol Cancer Ther, 2008, 7(8), 2348-2357.

Bruno RD, et al. Mol Cancer Ther, 2008, 7(9), 2828-2836.

Schayowitz A, et al. Mol Cancer Ther, 2008, 7(1), 121-132.

Schayowitz A, et al. Br J Cancer, 2010, 103(7), 12001-12007.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481